

The Recognition Theorem for $\text{Out}(F_n)$

Mark Feighn*

Michael Handel†

February 15, 2009

Abstract

Our goal is to find dynamic invariants that completely determine elements of the outer automorphism group $\text{Out}(F_n)$ of the free group F_n of rank n . To avoid finite order phenomena, we do this for *forward rotationless* elements. This is not a serious restriction. For example, there is $K_n > 0$ depending only on n such that, for all $\phi \in \text{Out}(F_n)$, ϕ^{K_n} is forward rotationless. An important part of our analysis is to show that rotationless elements are represented by particularly nice relative train track maps.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Marked Graphs and Outer Automorphisms	5
2.2	Paths, Circuits and Edge Paths	5
2.3	Automorphisms and Lifts	6
2.4	Lines and Laminations	7
2.5	Free Factor Systems	8
2.6	Relative Train Track Maps	9
2.7	Modifying relative train track maps	15
3	Forward Rotationless Outer Automorphisms	24
3.1	The Nielsen Approach to the Mapping Class Group	24
3.2	Principal Automorphisms	25
3.3	Rotationless Relative Train Track Maps and Principal Periodic Points	28
3.4	Rotationless is Rotationless	33
3.5	Properties of Forward Rotationless ϕ	35

*This material is based upon work supported by the National Science Foundation under Grant No. DMS-0805440.

†This material is based upon work supported by the National Science Foundation under Grant No. DMS-0706719 .

4	Completely Split Relative Train Track Maps	36
4.1	Definitions and Notation	37
4.2	Hard Splittings	39
4.3	CT versus IRT	41
4.4	A new move	47
4.5	Existence Theorem	48
4.6	Further properties of a CT	54
5	Recognition Theorem	58
5.1	The NEG Case	61
5.2	The EG Case	62

1 Introduction

The Thurston classification theorem for mapping class groups of surfaces inspired a surge in research on the outer automorphism group $\text{Out}(F_n)$ of the free group of rank n . One direction of this research is the development and use of relative train track maps which are the analog of the normal forms for mapping classes. Thurston's normal forms give rise to invariants that completely determine a mapping class, perhaps after passing to a finite power. In this paper we provide similar invariants for elements of $\text{Out}(F_n)$ and we add an important feature to relative train track maps.

We begin by recalling the invariants associated to a mapping class. After passing to a finite power, there is a decomposition of the surface into invariant subsurfaces on which the action of the mapping class is either a Dehn twist in an annulus, trivial or pseudo-Anosov; in the pseudo-Anosov case, each singular ray of the associated measured foliations is invariant. The mapping class is completely determined by the (isotopy classes of the) core curves of the annuli, the Dehn twist degrees, the pseudo-Anosov measured foliations and the expansion factors on the pseudo-Anosov measured foliations.

To see how this might generalize to $\text{Out}(F_n)$, consider the special case in which the mapping class ψ is a composition of disjoint Dehn twists and so is determined by the twisting circles and the degrees of twist. The dual point of view is useful here. The complementary components of the twisting curves are maximal subsurfaces on which ψ acts trivially. Viewing the mapping class group as the group of outer automorphism of the fundamental group $\pi_1(S)$ of the surface S , each complementary component determines, up to conjugacy, a subgroup of $\pi_1(S)$ of rank at least two that is the fixed subgroup $\text{Fix}(\Psi)$ of some automorphism Ψ of $\pi_1(S)$ that represents the outer automorphism ψ . If Ψ_1 and Ψ_2 are two distinct such automorphisms, corresponding to either the same or distinct subsurfaces, then $\text{Fix}(\Psi_1) \cap \text{Fix}(\Psi_2)$ is either trivial or a maximal cyclic group $\langle a \rangle$. In the latter case $\Psi_2 = i_a^d \Psi_1$ for some $d \neq 0$ where i_a is the inner automorphism determined by a ; moreover, the conjugacy class determined by a represents a twisting curve for μ with twisting degree $\pm d$.

This point of view, focusing on fixed subgroups of rank at least two and their intersections, is sufficient [6] if one restricts to elements of $\text{Out}(F_n)$ that have linear growth.¹ For general $\phi \in \text{Out}(F_n)$, we must also account for exponential and non-linear polynomial growth. To do this, we consider the boundary of the free group. An automorphism Φ induces a homeomorphism $\hat{\Phi}: \partial F_n \rightarrow \partial F_n$. In the linear case, the fixed point set $\text{Fix}(\hat{\Phi})$ of $\hat{\Phi}$ is equal to the boundary of the fixed subgroup $\text{Fix}(\Phi)$ and contains no isolated points. In the general case, however, there are isolated points, and these contain essential information about the automorphism. For example, the set of attracting laminations associated to ϕ can be recovered from the isolated attracting points.

The idea then is to replace $\text{Fix}(\Phi)$ with the set $\text{Fix}_N(\hat{\Phi})$ of non-isolated points and attractors in $\text{Fix}(\hat{\Phi})$; see [12], [14], [3] and [15] where this same idea has been used effectively. In section 3.2, we define the set $P(\phi) \subset \text{Aut}(F_n)$ of *principal automorphisms* representing ϕ . In the case of linear growth, Φ is principal if and only if $\text{Fix}(\Phi)$ has rank at least two. The invariants that determine ϕ , after possibly passing to a finite power, are the sets $\text{Fix}_N(\hat{\Phi})$ as Φ varies over $P(\phi)$, the expansion factors for the attracting laminations of ϕ and twisting coordinates associated to pairs of principal automorphisms whose fixed points sets intersect non-trivially.

It is common when studying elements $\phi \in \text{Out}(F_n)$ to ‘stabilize’ ϕ by replacing it with a power ϕ^k . In section 3, we specify a subset of $\text{Out}(F_n)$ whose elements require no stabilization. These outer automorphisms are said to be *forward rotationless*. In Lemma 4.43 we prove that there is $K_n > 0$, depending only on n , so that ϕ^{K_n} is forward rotationless for all $\phi \in \text{Out}(F_n)$. We also define what it means for a relative train track map $f: G \rightarrow G$ to be *rotationless* and prove (Proposition 3.29) that ϕ is forward rotationless if and only if some (every) relative train track map representing it is rotationless. (There is no need to add ‘forward’ to this terminology because f^k is only defined for $k \geq 1$.) It is easy to check if $f: G \rightarrow G$ is rotationless and if not to find the minimal k such that f^k is rotationless.

We can now state our main result. Complete details and further motivation are supplied in section 5. In addition to being of intrinsic interest this theorem is needed in [10]. The set of attracting laminations for ϕ is denoted $\mathcal{L}(\phi)$ and the expansion factor for ϕ on $\Lambda \in \mathcal{L}(\phi)$ is denoted $\text{PF}_\Lambda(\phi)$.

Theorem 5.3 (Recognition Theorem) *Suppose that $\phi, \psi \in \text{Out}(F_n)$ are forward rotationless and that*

- (1) $\text{PF}_\Lambda(\phi) = \text{PF}_\Lambda(\psi)$, for all $\Lambda \in \mathcal{L}(\phi) = \mathcal{L}(\psi)$; and
- (2) *there is bijection $B: P(\phi) \rightarrow P(\psi)$ such that:*

- (i) **(fixed sets preserved)** $\text{Fix}_N(\hat{\Phi}) = \text{Fix}_N(\widehat{B(\Phi)})$; and

¹ $\phi \in \text{Out}(F_n)$ has *linear growth* if, for all conjugacy classes of elements $a \in F_n$, the cyclically reduced word length of the conjugacy class of $\phi^m(a)$ is bounded by a linear function (depending on a) of m .

(ii) (**twist coordinates preserved**) if $w \in \text{Fix}(\Phi)$ and $\Phi, i_w \Phi \in P(\phi)$, then $B(i_w \Phi) = i_w B(\Phi)$.

Then $\phi = \psi$.

In the course of proving Theorem 5.3, we construct relative train track maps that are better than the those constructed in [2]. We also reorganize elements of the theory to make future modification and referencing of results easier.

The idea behind relative train track maps is that one can study the action of an outer automorphism ϕ on conjugacy classes in F_n or on ∂F_n by analyzing the action of a homotopy equivalence $f : G \rightarrow G$ of a marked graph G representing ϕ on paths, circuits and lines in G . For simplicity, suppose that σ is a finite path in G , which by convention is always assumed to be the immersed image of a compact interval. The image $f(\sigma)$ of σ is homotopic rel endpoints to a path denoted $f_{\#}(\sigma)$. Replacing $f(\sigma)$ with $f_{\#}(\sigma)$ is called *tightening* and is analogous to replacing a word in a set of generators for F_n with a reduced word in those generators. A decomposition of σ into subpaths $\sigma = \sigma_1 \dots \sigma_m$ is a *splitting* if $f_{\#}^k(\sigma) = f_{\#}^k(\sigma_1) \dots f_{\#}^k(\sigma_m)$ for all $k \geq 0$; i.e. if one can tighten the image of σ under any iterate of f by tightening the images of the σ_i 's. The more one can split σ and the better one can understand the subpaths σ_i , the more effectively one can analyze the iterates $f_{\#}^k(\sigma)$.

Relative train track maps were defined and constructed in [4] with exponentially growing strata in mind. Few restrictions were placed on the non-exponentially growing strata. This was rectified in [2] where *improved relative train tracks (IRTs)* are defined and shown to exist for a sufficiently high, but unspecified, iterate of ϕ . For our current application, IRTs are inadequate. In Theorem 4.29, we prove that every forward rotationless ϕ is represented by a relative train track map $f : G \rightarrow G$ that has all the essential properties of an IRT (see section 4.3) and has the additional feature that, for all σ and all sufficiently large k , there is a canonical splitting (called the *complete splitting*) of $f_{\#}^k(\sigma)$ into simple, explicitly described subpaths. (Splittings in an IRT are not canonical and the subpaths σ_i are understood more inductively than explicitly.) Such $f : G \rightarrow G$, called *CTs*, are used in the proof of the Recognition Theorem and in the classification of abelian subgroups given in [10]. It is likely that the existence of complete splittings will be useful in other contexts as well. For example, complete splittings are *hard splittings* as defined in [5].

2 Preliminaries

Fix $n \geq 2$ and let F_n be the free group of rank n . Denote the automorphism group of F_n by $\text{Aut}(F_n)$, the group of inner automorphisms of F_n by $\text{Inn}(F_n)$ and the group of outer automorphisms of F_n by $\text{Out}(F_n) = \text{Aut}(F_n)/\text{Inn}(F_n)$. We follow the convention that elements of $\text{Aut}(F_n)$ are denoted by upper case Greek letters and that the same Greek letter in lower case denotes the corresponding element of $\text{Out}(F_n)$. Thus $\Phi \in$

$\text{Aut}(F_n)$ represents $\phi \in \text{Out}(F_n)$.

2.1 Marked Graphs and Outer Automorphisms

Identify F_n with $\pi_1(R_n, *)$ where R_n is the rose with one vertex $*$ and n edges. A *marked graph* G is a graph of rank n , all of whose vertices have valence at least two, equipped with a homotopy equivalence $m : R_n \rightarrow G$ called a *marking*. Letting $b = m(*) \in G$, the marking determines an identification of F_n with $\pi_1(G, b)$. It is often assumed that G does not have valence two vertices, but such vertices occur naturally in relative train track theory so we allow them.

A homotopy equivalence $f : G \rightarrow G$ and a path σ from b to $f(b)$ determines an automorphism of $\pi_1(G, b)$ and hence an element of $\text{Aut}(F_n)$. If f fixes b and no path is specified, then we use the trivial path. This construction depends only on the homotopy class of σ and, as the homotopy class varies, the automorphism ranges over all representatives of the associated outer automorphism ϕ . We say that $f : G \rightarrow G$ *represents* ϕ . We always assume that f maps vertices to vertices and that the restriction of f to any edge is an immersion.

2.2 Paths, Circuits and Edge Paths

Let Γ be the universal cover of a marked graph G and let $pr : \Gamma \rightarrow G$ be the covering projection. A proper map $\tilde{\sigma} : J \rightarrow \Gamma$ with domain a (possibly infinite) closed interval J will be called a *path in* Γ if it is an embedding or if J is finite and the image is a single point; in the latter case we say that $\tilde{\sigma}$ is a *trivial path*. If J is finite, then $\tilde{\sigma} : J \rightarrow \Gamma$ is homotopic rel endpoints to a unique (possibly trivial) path $[\tilde{\sigma}]$; we say that $[\tilde{\sigma}]$ *is obtained from* $\tilde{\sigma}$ *by tightening*. If $\tilde{f} : \Gamma \rightarrow \Gamma$ is a lift of a homotopy equivalence $f : G \rightarrow G$, we denote $[\tilde{f}(\tilde{\sigma})]$ by $\tilde{f}_\#(\tilde{\sigma})$.

We will not distinguish between paths in Γ that differ only by an orientation preserving change of parametrization. Thus we are interested in the oriented image of $\tilde{\sigma}$ and not $\tilde{\sigma}$ itself. If the domain of $\tilde{\sigma}$ is finite, then the image of $\tilde{\sigma}$ has a natural decomposition as a concatenation $\tilde{E}_1\tilde{E}_2\ldots\tilde{E}_{k-1}\tilde{E}_k$ where $\tilde{E}_i$, $1 < i < k$, is an edge of Γ , $\tilde{E}_1$ is the terminal segment of an edge and $\tilde{E}_k$ is the initial segment of an edge. If the endpoints of the image of $\tilde{\sigma}$ are vertices, then $\tilde{E}_1$ and $\tilde{E}_k$ are full edges. The sequence $\tilde{E}_1\tilde{E}_2\ldots\tilde{E}_k$ is called *the edge path associated to* $\tilde{\sigma}$. This notation extends naturally to the case that the interval of domain is half-infinite or bi-infinite. In the former case, an edge path has the form $\tilde{E}_1\tilde{E}_2\ldots$ or $\ldots\tilde{E}_{-2}\tilde{E}_{-1}$ and in the latter case has the form $\ldots\tilde{E}_{-1}\tilde{E}_0\tilde{E}_1\tilde{E}_2\ldots$.

A *path in* G is the composition of the projection map pr with a path in Γ . Thus a map $\sigma : J \rightarrow G$ with domain a (possibly infinite) closed interval will be called a *path* if it is an immersion or if J is finite and the image is a single point; paths of the latter type are said to be *trivial*. If J is finite, then $\sigma : J \rightarrow G$ is homotopic rel endpoints to a unique (possibly trivial) path $[\sigma]$; we say that $[\sigma]$ *is obtained from* σ

by *tightening*. For any lift $\tilde{\sigma} : J \rightarrow \Gamma$ of σ , $[\sigma] = pr[\tilde{\sigma}]$. We denote $[f(\sigma)]$ by $f_{\#}(\sigma)$. We do not distinguish between paths in G that differ by an orientation preserving change of parametrization. The *edge path associated to σ* is the projected image of the edge path associated to a lift $\tilde{\sigma}$. Thus the edge path associated to a path with finite domain has the form $E_1 E_2 \dots E_{k-1} E_k$ where E_i , $1 < i < k$, is an edge of G , E_1 is the terminal segment of an edge and E_k is the initial segment of an edge. We will identify paths with their associated edge paths whenever it is convenient.

We reserve the word *circuit* for an immersion $\sigma : S^1 \rightarrow G$. Any homotopically non-trivial map $\sigma : S^1 \rightarrow G$ is homotopic to a unique circuit $[\sigma]$. As was the case with paths, we do not distinguish between circuits that differ only by an orientation preserving change in parametrization and we identify a circuit σ with a *cyclically ordered edge path* $E_1 E_2 \dots E_k$.

A path or circuit *crosses* or *contains* an edge if that edge occurs in the associated edge path. For any path σ in G define $\bar{\sigma}$ to be ‘ σ with its orientation reversed’. For notational simplicity, we sometimes refer to the inverse of $\tilde{\sigma}$ by $\tilde{\sigma}^{-1}$.

A decomposition of a path or circuit into subpaths is a *splitting* for $f : G \rightarrow G$ and is denoted $\sigma = \dots \cdot \sigma_1 \cdot \sigma_2 \cdot \dots$ if $f_{\#}^k(\sigma) = \dots f_{\#}^k(\sigma_1) f_{\#}^k(\sigma_2) \dots$ for all $k \geq 0$. In other words, a decomposition of σ into subpaths σ_i is a splitting if one can tighten the image of σ under any iterate of $f_{\#}$ by tightening the images of the σ_i ’s.

A path σ is a *periodic Nielsen path* if σ is non-trivial and $f_{\#}^k(\sigma) = \sigma$ for some $k \geq 1$. The minimal such k is the *period* of σ and if the period is one then σ is a *Nielsen path*. A (periodic) Nielsen path is *indivisible* if it does not decompose as a concatenation of non-trivial (periodic) Nielsen subpaths. A path or circuit is *root-free* if it is not equal to μ^k for some simpler path μ and some $k > 1$.

2.3 Automorphisms and Lifts

Section 1 of [12] and section 2.1 of [3] are good sources for facts that we record below without specific references. The universal cover Γ of a marked graph G with marking $m : R_n \rightarrow G$ is a simplicial tree. We always assume that a base point $\tilde{b} \in \Gamma$ projecting to $b = m(*) \in G$ has been chosen, thereby defining an action of F_n on Γ . The set of ends $\mathcal{E}(\Gamma)$ of Γ is naturally identified with the boundary ∂F_n of F_n and we make implicit use of this identification throughout the paper.

Each non-trivial $c \in F_n$ acts by a *covering translation* $T_c : \Gamma \rightarrow \Gamma$ and each T_c induces a homeomorphism $\hat{T}_c : \partial F_n \rightarrow \partial F_n$ that fixes two points, a sink T_c^+ and a source T_c^- . The line in Γ whose ends converge to T_c^- and T_c^+ is called the *axis* of T_c and is denoted A_c . The image of A_c in G is the circuit corresponding to the conjugacy class of c .

If $f : G \rightarrow G$ represents $\phi \in \text{Out}(F_n)$ then a path σ from b to $f(b)$ determines both an automorphism representing ϕ and a lift of f to Γ . This defines a bijection between the set of lifts $\tilde{f} : \Gamma \rightarrow \Gamma$ of $f : G \rightarrow G$ and the set of automorphisms $\Phi : F_n \rightarrow F_n$ representing ϕ . Equivalently, this bijection is defined by $\tilde{f} T_c = T_{\Phi(c)} \tilde{f}$ for all $c \in F_n$.

We say that $\tilde{f}$ corresponds to Φ or is determined by Φ and vice versa. Under the identification of $\mathcal{E}(\Gamma)$ with ∂F_n , a lift $\tilde{f}$ determines a homeomorphism $\hat{f}$ of ∂F_n . An automorphism Φ also determines a homeomorphism $\hat{\Phi}$ of ∂F_n and $\hat{f} = \hat{\Phi}$ if and only if $\tilde{f}$ corresponds to Φ . In particular, $\hat{i}_c = \hat{T}_c$ for all $c \in F_n$ where $i_c(w) = cwc^{-1}$ is the inner automorphism of F_n determined by c . We use the notation $\hat{f}$ and $\hat{\Phi}$ interchangeably depending on the context.

We are particularly interested in the dynamics of $\hat{f} = \hat{\Phi}$. The following two lemmas are contained in Lemmas 2.3 and 2.4 of [3] and in Proposition 1.1 of [12].

Lemma 2.1. *Assume that $\tilde{f} : \Gamma \rightarrow \Gamma$ corresponds to $\Phi \in \text{Aut}(F_n)$. Then the following are equivalent:*

- (i) $c \in \text{Fix}(\Phi)$.
- (ii) T_c commutes with $\tilde{f}$.
- (iii) $\hat{T}_c$ commutes with $\hat{f}$.
- (iv) $\text{Fix}(\hat{T}_c) \subset \text{Fix}(\hat{f}) = \text{Fix}(\hat{\Phi})$.
- (v) $\text{Fix}(\hat{f}) = \text{Fix}(\hat{\Phi})$ is $\hat{T}_c$ -invariant.

Remark 2.2. It is not hard to see that $T_c^+ \in \text{Fix}(\Phi)$ if and only if $T_c^- \in \text{Fix}(\Phi)$.

A point $P \in \partial F_n$ is an *attractor* for $\hat{\Phi}$ if it has a neighborhood U such that $\hat{\Phi}(U) \subset U$ and such that $\bigcap_{n=1}^{\infty} \hat{\Phi}^n(U) = P$. If Q is an attractor for $\hat{\Phi}^{-1}$ then we say that it is a *repeller* for $\hat{\Phi}$.

Lemma 2.3. *Assume that $\tilde{f} : \Gamma \rightarrow \Gamma$ corresponds to $\Phi \in \text{Aut}(F_n)$ and that $\text{Fix}(\hat{\Phi}) \subset \partial F_n$ contains at least three points. Denote $\text{Fix}(\hat{\Phi})$ by $\mathbb{F}$ and the corresponding subgroup of covering translations of Γ by $\mathbb{T}(\Phi)$. Then*

- (i) $\partial \mathbb{F}$ is naturally identified with the closure of $\{T_c^{\pm} : T_c \in \mathbb{T}(\Phi)\}$ in ∂F_n . None of these points is isolated in $\text{Fix}(\hat{\Phi})$.
- (ii) Each point in $\text{Fix}(\hat{\Phi}) \setminus \partial \mathbb{F}$ is isolated and is either an attractor or a repeller for the action of $\hat{\Phi}$.
- (iii) There are only finitely many $\mathbb{T}(\Phi)$ -orbits in $\text{Fix}(\hat{\Phi}) \setminus \partial \mathbb{F}$.

2.4 Lines and Laminations

Suppose that Γ is the universal cover of a marked graph G . An unoriented bi-infinite path in Γ is called a *line* in Γ . The *space of lines* in Γ is denoted $\tilde{\mathcal{B}}(\Gamma)$ and is equipped with what amounts to the compact-open topology. Namely, for any finite path $\tilde{\alpha}_0 \subset \Gamma$ (with endpoints at vertices if desired), define $N(\tilde{\alpha}_0) \subset \tilde{\mathcal{B}}(\Gamma)$ to be the set of lines in

Γ that contain $\tilde{\alpha}_0$ as a subpath. The sets $N(\tilde{\alpha}_0)$ define a basis for the topology on $\tilde{\mathcal{B}}(\Gamma)$.

An unoriented bi-infinite path in G is called a *line in G* . The space of lines in G is denoted $\mathcal{B}(G)$. There is a natural projection map from $\tilde{\mathcal{B}}(\Gamma)$ to $\mathcal{B}(G)$ and we equip $\mathcal{B}(G)$ with the quotient topology.

A line in Γ is determined by the unordered pair of its endpoints (P, Q) and so corresponds to a point in the *space of abstract lines* defined to be $\tilde{\mathcal{B}} := ((\partial F_n \times \partial F_n) \setminus \Delta) / \mathbb{Z}_2$, where Δ is the diagonal and where $\mathbb{Z}_2$ acts on $\partial F_n \times \partial F_n$ by interchanging the factors. The action of F_n on ∂F_n induces an action of F_n on $\tilde{\mathcal{B}}$ whose quotient space is denoted $\mathcal{B}$. The ‘endpoint map’ defines a homeomorphism between $\tilde{\mathcal{B}}$ and $\tilde{\mathcal{B}}(\Gamma)$ and we use this implicitly to identify $\tilde{\mathcal{B}}$ with $\tilde{\mathcal{B}}(\Gamma)$ and hence $\tilde{\mathcal{B}}(\Gamma)$ with $\tilde{\mathcal{B}}(\Gamma')$ where Γ' is the universal cover of any other marked graph G' . There is a similar identification of $\mathcal{B}(G)$ with $\mathcal{B}$ and with $\mathcal{B}(G')$. We sometimes say that the line in G or Γ corresponding to an abstract line is *the realization* of that abstract line in G or Γ .

A closed set of lines in G or a closed F_n -invariant set of lines in Γ is called a *lamination* and the lines that compose it are called *leaves*. If Λ is a lamination in G then we denote its pre-image in Γ by $\tilde{\Lambda}$ and vice-versa.

Suppose that $f : G \rightarrow G$ represents ϕ and that $\tilde{f}$ is a lift of f . If $\tilde{\gamma}$ is a line in Γ with endpoints P and Q , then there is a bounded homotopy from $\tilde{f}(\tilde{\gamma})$ to the line $\tilde{f}_\#(\tilde{\gamma})$ with endpoints $\tilde{f}(P)$ and $\tilde{f}(Q)$. This defines an action $\tilde{f}_\#$ of $\tilde{f}$ on lines in Γ . If $\Phi \in \text{Aut}(F_n)$ corresponds to $\tilde{f}$ then $\Phi_\# = \tilde{f}_\#$ is described on abstract lines by $(P, Q) \mapsto (\hat{\Phi}(P), \hat{\Phi}(Q))$. There is an induced action $\phi_\#$ of ϕ on lines in G and in particular on laminations in G .

To each $\phi \in \text{Out}(F_n)$ is associated a finite ϕ -invariant set of laminations $\mathcal{L}(\phi)$ called the set of *attracting laminations* for ϕ . The individual laminations need not be ϕ -invariant. By definition (see Definition 3.1.5 of [2]) $\mathcal{L}(\phi) = \mathcal{L}(\phi^k)$ for all $k \geq 1$ and each $\Lambda \in \mathcal{L}(\phi)$ contains birecurrent leaves, called *generic leaves*, whose weak closure is all of Λ . Complete details on $\mathcal{L}(\phi)$ can be found in section 3 of [2].

A point $P \in \partial F_n$ determines a lamination $\Lambda(P)$, called *the accumulation set of P* , as follows. Let Γ be the universal cover of a marked graph G and let $\tilde{R}$ be any ray in Γ converging to P . A line $\tilde{\sigma} \subset \Gamma$ belongs to $\tilde{\Lambda}(P)$ if every finite subpath of $\tilde{\sigma}$ is contained in some translate of $\tilde{R}$. Since any two rays converging to P have a common infinite end, this definition is independent of the choice of $\tilde{R}$. The bounded cancellation lemma implies (cf. Lemma 3.1.4 of [2]) that this definition is independent of the choice of G and Γ and that $\hat{\Phi}_\#(\tilde{\Lambda}(P)) = \tilde{\Lambda}(\hat{\Phi}(P))$. In particular, if $P \in \text{Fix}(\hat{\Phi})$ then $\Lambda(P)$ is $\phi_\#$ -invariant.

2.5 Free Factor Systems

The conjugacy class of a free factor F^i of F_n is denoted $[[F^i]]$. If $F^1, \dots, F^k$ are non-trivial free factors and if $F^1 * \dots * F^k$ is a free factor then we say that the

collection $\{[[F^1]], \dots, [[F^k]]\}$ is a *free factor system*. For example, if G is a marked graph and $G_r \subset G$ is a subgraph with non-contractible components $C_1, \dots, C_k$ then the conjugacy class $[[\pi_1(C_i)]]$ of the fundamental group of C_i is well defined and the collection of these conjugacy classes is a free factor system denoted $\mathcal{F}(G_r)$; we say that G_r *realizes* $\mathcal{F}(G_r)$.

The image of a free factor F under an element of $\mathbf{Aut}(F_n)$ is a free factor. This induces an action of $\mathbf{Out}(F_n)$ on the set of free factor systems. We sometimes say that a free factor is ϕ -invariant when we really mean that its conjugacy class is ϕ -invariant. If $[[F]]$ is ϕ -invariant then F is Φ -invariant for some automorphism Φ representing ϕ and $\Phi|F$ determines a well defined element $\phi|F$ of $\mathbf{Out}(F)$.

The conjugacy class $[a]$ of $a \in F_n$ is *carried by* $[[F^i]]$ if F^i contains a representative of $[a]$. Sometimes we say that a is carried by F^i when we really mean that $[a]$ is carried by $[[F^i]]$. If G is a marked graph and G_r is a subgraph of G such that $[[F^i]] = [[\pi_1(G_r)]]$, then $[a]$ is carried by $[[F^i]]$ if and only if the circuit in G that represents $[a]$ is contained in G_r . We say that *an abstract line ℓ is carried by $[[F^i]]$* if its realization in G is contained in G_r for some, and hence any, G and G_r as above. Equivalently, ℓ is the limit of periodic lines corresponding to $[c_i]$ where each $[c_i]$ is carried by $[[F^i]]$. A collection W of abstract lines and conjugacy classes in F_n is carried by a free factor system $\mathcal{F} = \{[[F^1]], \dots, [[F^k]]\}$ if each element of W is carried by some F^i .

There is a partial order $\sqsubset$ on free factor systems generated by inclusion. More precisely, $[[F^1]] \sqsubset [[F^2]]$ if F^1 is conjugate to a free factor of F^2 and $\mathcal{F}_1 \sqsubset \mathcal{F}_2$ if for each $[[F^i]] \in \mathcal{F}_1$ there exists $[[F^j]] \in \mathcal{F}_2$ such that $[[F^i]] \sqsubset [[F^j]]$.

The *complexity* of a free factor system is defined on page 531 of [2]. Rather than repeat the definition, we recall the three properties of complexity that we use. The first is that if $\mathcal{F}_1 \sqsubset \mathcal{F}_2$ for distinct free factor systems $\mathcal{F}_1$ and $\mathcal{F}_2$ then the complexity of $\mathcal{F}_1$ is less than the complexity of $\mathcal{F}_2$. This is immediate from the definition. The second is Corollary 2.6.5. of [2].

Lemma 2.4. *For any collection W of abstract lines there is a unique free factor system $\mathcal{F}(W)$ of minimal complexity that carries every element of W . If W is a single element then $\mathcal{F}(W)$ has a single element.*

The third is an immediate consequence of the uniqueness of $\mathcal{F}(W)$.

Corollary 2.5. *If a collection W of abstract lines and conjugacy classes in F_n is ϕ -invariant then $\mathcal{F}(W)$ is ϕ -invariant.*

Further details on free factor systems can be found in section 2.6 of [2].

2.6 Relative Train Track Maps

In this section we review and set notation for relative train track maps as defined in [4].

Suppose that G is a marked graph and that $f : G \rightarrow G$ is a homotopy equivalence representing $\phi \in \text{Out}(F_n)$. A *filtration* of G is an increasing sequence $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ of subgraphs, each of whose components contains at least one edge. If $f(G_i) \subset G_i$ for all i then we say that $f : G \rightarrow G$ *respects the filtration* or that the *filtration is f -invariant*. A path or circuit has *height* r if it is contained in G_r but not G_{r-1} . A lamination has height r if each leaf in its realization in G has height at most r and some leaf has height r .

The r^{th} *stratum* H_r is defined to be the closure of $G_r \setminus G_{r-1}$. To each stratum H_r there is an associated square matrix M_r , called the *transition matrix for H_r* , whose ij^{th} entry is the number of times that the i^{th} edge (in some ordering of the edges of H_r) crosses the j^{th} edge in either direction. By enlarging the filtration, we may assume that each M_r is either irreducible or the zero matrix. We say that H_r is an *irreducible stratum* if M_r is irreducible and is a *zero stratum* if M_r is the zero matrix.

If M_r is irreducible and the Perron-Frobenius eigenvalue of M_r is 1, then M_r is a permutation matrix and H_r is *non-exponentially growing* or simply *NEG*. After subdividing and replacing the given NEG stratum with a pair of NEG strata if necessary, the edges $\{E_1, \dots, E_l\}$ of H_r can be oriented and ordered so that $f(E_i) = E_{i+1}u_i$ where $u_i \subset G_{r-1}$ and where indices are taken mod l . We always assume that edges in an NEG stratum have been so oriented and ordered. If each u_i is trivial then $f^l(E_i) = E_i$ for all i and we say that $E_i [H_r]$ is a *periodic edge [stratum]* with period l or a *fixed edge [stratum]* if $l = 1$. If each u_i is a Nielsen path then the combinatorial length of $f_{\#}^k(E_i)$ is bounded by a linear function of k for all i and we say that $E_i [H_r]$ is a *linear edge [stratum]*.

If M_r is irreducible and if the Perron-Frobenius eigenvalue of M_r is greater than 1 then H_r is an *exponentially growing stratum* or simply an *EG stratum*. If H_r is EG and $\alpha \subset G_{r-1}$ is a non-trivial path with endpoints in $H_r \cap G_{r-1}$ then we say that α is a *connecting path for H_r* . If H_r is EG and σ is a path with height r then we sometimes say that σ *has EG height*.

A *direction* d at $x \in G$ is the germ of an initial segment of an oriented edge (or partial edge if x is not a vertex) based at x . There is an f -induced map Df on directions and we say that d is a *periodic direction* if it is periodic under the action of Df ; if the period is one then d is a *fixed direction*. Thus, the direction determined by an oriented edge E is fixed if and only if E is the initial edge of $f(E)$. Two directions with the same basepoint belong to the same *gate* if they are identified by some iterate of Df . If x is a periodic point then the number of gates based at x is equal to the number of periodic directions based at x .

A *turn* is an unordered pair of directions with a common base point. The turn is *nondegenerate* if it is defined by distinct directions and is *degenerate* otherwise. If $E_1 E_2 \dots E_{k-1} E_k$ is the edge path associated to a path σ , then we say that σ *contains the turns* $(\bar{E}_i, E_{i+1})$ for $1 \leq i \leq k-1$. A turn is *illegal* with respect to $f : G \rightarrow G$ if its image under some iterate of Df is degenerate; a turn is *legal* if it is not illegal. Equivalently, a turn is legal if and only if it is defined by directions that belong to

distinct gates. A *path or circuit* $\sigma \subset G$ is *legal* if it contains only legal turns. If $\sigma \subset G_r$ does not contain any illegal turns in H_r , meaning that both directions correspond to edges of H_r , then σ is *r-legal*. It is immediate from the definitions that Df maps legal turns to legal turns and that the restriction of f to a legal path is an immersion.

We recall the definition of relative train track map from page 38 of [4].

Definition 2.6. A homotopy equivalence $f : G \rightarrow G$ representing ϕ is a *relative train track map* if it satisfies the following conditions for every EG stratum H_r of an f -invariant filtration $\mathcal{F}$.

- (RTT-i) Df maps the set of directions in H_r with basepoints at vertices to itself; in particular every turn with one direction in H_r and the other in G_{r-1} is legal.
- (RTT-ii) If α is a connecting path for H_r then $f_{\#}(\alpha)$ is a connecting path for H_r ; in particular, $f_{\#}(\alpha)$ is nontrivial.
- (RTT-iii) If $\alpha \subset G_r$ is *r-legal* then $f_{\#}(\alpha)$ is *r-legal*.

Remark 2.7. If $f : G \rightarrow G$ is a relative train track map, then so is f^k for $k > 0$.

A subgraph C of G is *wandering* if $f^k(C)$ is contained in the closure of $G \setminus J$ for all $k > 0$; otherwise C is *non-wandering*. Each edge in a wandering subgraph is contained in a zero stratum. If C is a component of a filtration element then C is non-wandering if and only if $f^i(C) \subset C$ for some $i > 0$.

Remark 2.8. Suppose that H_r is an EG stratum and that $f : G \rightarrow G$ satisfies (RTT-i) for H_r . Then $f(H_r \cap G_{r-1}) \subset H_r \cap G_{r-1}$ and verifying (RTT-ii) for H_r reduces to showing that $f_{\#}(\alpha)$ is nontrivial for each connecting path α for H_r . Suppose that the component C of G_{r-1} that contains α is non-wandering. In checking (RTT-ii) there is no loss in replacing f by f^i so we may assume that $f(C) \subset C$. If f permutes the elements of the finite set $H_r \cap C$, then $f_{\#}(\alpha)$ is nontrivial for each $\alpha \subset C$. If f identifies two of these points then they can be connected by an arc $\alpha \subset C$ such that $f_{\#}(\alpha)$ is trivial (because $f|_C : C \rightarrow C$ is a homotopy equivalence). This proves that if C is a non-wandering component of G_{r-1} then (RTT-ii) holds for all $\alpha \subset C$ if and only if $H_r \cap C \subset \text{Per}(f)$.

The most common applications of the relative train track properties are contained in following lemma.

Lemma 2.9. *Suppose that $f : G \rightarrow G$ is a relative train track map and that H_r is an EG stratum.*

- (1) *Suppose that a vertex v of H_r is contained in a component C of G_{r-1} that is non-contractible or more generally satisfies $f^i(C) \subset C$ for some $i > 0$. Then v is periodic and has at least one periodic direction in H_r .*

- (2) If σ is an r -legal circuit or path of height r with endpoints, if any, at vertices of H_r then the decomposition of σ into single edges in H_r and maximal subpaths in G_{r-1} is a splitting.

Proof. The first item follows from Remark 2.8. The second item is contained in Lemma 5.8 of [4]. $\square$

Lemma 2.10. *If H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ and $x \in H_r$ is either a vertex or a periodic point then there is a legal turn in G_r that is based at x . In particular, there are at least two gates in G_r that are based at x .*

Proof. There exists $j > 0$ and a point y in the interior of an edge E of H_r so that $x = f^j(y)$. By (RTT-iii), x is in the interior of an r -legal path. Moreover the turn at x determined by this path is legal by Properties (RTT-iii) and (RTT-i). $\square$

The following lemma describes indivisible periodic Nielsen paths with EG height.

Lemma 2.11. *Suppose that $f : G \rightarrow G$ is a relative train track map and that H_r is an EG stratum.*

- (1) *There are only finitely many indivisible periodic Nielsen paths of height r .*
- (2) *If σ is an indivisible periodic Nielsen path of height r then $\sigma = \alpha\beta$ where α and β are r -legal paths that begin and end with directions in H_r and the turn $(\bar{\alpha}, \beta)$ is illegal. Moreover, if $\bar{\alpha}(k)$ and $\beta(k)$ are the initial segments of $\bar{\alpha}$ and β that are identified by $f_{\#}^k$ then the $\bar{\alpha}(k)\beta(k)$'s are an increasing sequence of subpaths whose union is the interior of σ .*
- (3) *An indivisible periodic Nielsen path σ of height r has period 1 if and only if the initial and terminal directions of σ are fixed.*

Proof. Item (1) and the first part of (2) is contained in the statement of Lemma 5.11 of [4]. The moreover part of (2) is contained in the proof of that lemma. We now turn our attention to (3).

Suppose that σ is an indivisible periodic Nielsen path of height r and period p and that $\sigma = \alpha\beta$ as in (2). After subdividing at the endpoints of $f_{\#}^j(\sigma)$ for $0 \leq j \leq p-1$ (which clearly preserves the property of being a relative train track map) we may assume that the endpoints of σ are vertices. Since α and $\bar{\beta}$ are r -legal and begin with edges in H_r , Lemma 2.9(2) implies that Df maps the initial directions of α and $\bar{\beta}$ to the initial direction of $f_{\#}(\alpha)$ and $f_{\#}(\bar{\beta})$ respectively. The only if part of (3) therefore follows from the fact that $f_{\#}(\sigma)$ is obtained from $f_{\#}(\alpha)$ and $f_{\#}(\bar{\beta})$ by cancelling their maximal common terminal segment.

Assume now that the first edge E of α determines a fixed direction and write $f_{\#}(\sigma) = \alpha_1\beta_1$ as in (2). Then E is also the first edge in α_1 . The moreover part of (2) implies that both α and α_1 are initial segments of $f_{\#}^{Np}(E)$ for all sufficiently large N . Thus either α is an initial segment of α_1 or α_1 is an initial segment of α . The

same argument applies to $\bar{\beta}$ and $\bar{\beta}_1$. The difference between the number of H_r edges in $f_{\#}^{Np}(\alpha)$ and the number of H_r edges in $f_{\#}^{Np}(\bar{\beta})$ is independent of N and similarly for α_1 and $\bar{\beta}_1$. It follows that either $\alpha \subset \alpha_1$ and $\bar{\beta} \subset \bar{\beta}_1$ or $\alpha_1 \subset \alpha$ and $\bar{\beta}_1 \subset \bar{\beta}$. For concreteness assume the former.

We claim that $\alpha_1 = \alpha$. If not then $\alpha_1 = \alpha\gamma$ for some non-trivial γ . The path $\alpha_2 := \bar{\beta}\gamma$ is a subpath of $f_{\#}^{Np}(\bar{\beta})$ for large N and so is r -legal. The path $\alpha_2\beta_1 = [(\bar{\beta}\bar{\alpha})(\alpha_1\beta_1)]$ is a non-trivial periodic Nielsen path with exactly one illegal turn in H_r and is therefore indivisible. By construction, α_2 and $\bar{\beta}$, and hence α_2 and $\bar{\beta}_1$, have a non-trivial common initial subpath in H_r . This implies as above that α_2 and $\bar{\beta}_1$ are initial subpaths of a common path δ . They cannot be equal so one is a proper initial subpath of the other. But then the difference between the number of H_r edges in $f_{\#}^{Np}(\alpha_2)$ and the number of H_r edges in $f_{\#}^{Np}(\bar{\beta}_1)$ grows exponentially in N in contradiction to the fact that $\alpha_2\beta_1$ is a periodic Nielsen path. This contradiction verifies the claim that $\alpha = \alpha_1$. The symmetric argument implies that $\beta = \beta_1$ so $p = 1$. $\square$

We extend Lemma 2.11(1) as follows.

Lemma 2.12. *For any relative train track map $f : G \rightarrow G$ there are only finitely many points that are endpoints of an indivisible periodic Nielsen path σ .*

Proof. Let S_r be the set of endpoints of indivisible periodic Nielsen paths σ with height at most r . We prove that S_r is finite for all r by induction on r . Since S_0 is empty we assume that S_{r-1} is finite and prove that S_r is finite. This is obvious if H_r is a zero stratum (because $S_r = S_{r-1}$) and follows from Lemma 2.11(1) if H_r is EG. We may therefore assume that H_r is NEG and that there is an indivisible Nielsen path σ of height r . Lemma 4.1.4 of [2] implies that H_r is not periodic (because σ is indivisible) and that edges of σ , other than perhaps the first and last, are not contained in H_r . The only periodic points in H_r are vertices so we may assume that $\sigma = Eu$ where E is an edge of H_r and $u \subset G_{r-1}$. If $\sigma' = Eu'$ is another such indivisible periodic Nielsen path then $\bar{u}u'$ is an indivisible periodic Nielsen path with height less than r and so its endpoints are contained in S_{r-1} . Thus S_r is obtained from S_{r-1} by adding at most two points for each edge in H_r . This completes the inductive step. $\square$

An irreducible matrix M is *aperiodic* if M^k has all positive entries for some $k \geq 1$. For example, if some diagonal element of M is non-zero then M is aperiodic. If H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ and if the transition matrix M_r is aperiodic, then H_r is said to be an *aperiodic EG stratum*. For each aperiodic EG stratum there is a unique ϕ -invariant attracting lamination $\Lambda_r \in \mathcal{L}(\phi)$ of height r . If every EG stratum is aperiodic then every element of $\mathcal{L}(\phi)$ is related to an EG stratum in this way. See Definition 3.1.12 of [2] and the surrounding material for details.

The following lemma produces rays and lines associated to aperiodic EG strata of a relative train track map.

Lemma 2.13. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$, that $\tilde{f} : \Gamma \rightarrow \Gamma$ is a lift and that $\tilde{v} \in \text{Fix}(\tilde{f})$.*

- (1) *If E is an oriented edge in H_r and $\tilde{E}$ is a lift that determines a fixed direction at $\tilde{v}$, then there is an r -legal ray $\tilde{R} \subset \Gamma$ that begins with $\tilde{E}$, intersects $\text{Fix}(\tilde{f})$ only in $\tilde{v}$ and that converges to an attractor $P \in \text{Fix}(\hat{f})$ whose accumulation set is the (necessarily ϕ -invariant) element Λ_r of $\mathcal{L}(\phi)$ whose realization in G has height r .*
- (2) *Suppose that $E' \neq E$ is another oriented edge in H_r , that $\tilde{E}'$ determines a fixed direction at $\tilde{v}$ and that $\tilde{R}'$ is the ray associated to $\tilde{E}'$ as in (1). Suppose further that the turn $(\tilde{E}, \tilde{E}')$ is contained in the path $f_{\#}^k(E'')$ for some $k \geq 1$ and some edge E'' of H_r . Then the line $\tilde{R}^{-1}\tilde{R}'$ is a generic leaf of $\tilde{\Lambda}_r$.*

Proof. Lemma 2.9(2) and (RTT-i) imply that $\tilde{f}(\tilde{E}) = \tilde{E} \cdot \tilde{\mu}_1$ for some non-trivial r -legal subpath $\tilde{\mu}_1$ of height r that ends with an edge of H_r . Applying Lemma 2.9(2) again, we have $\tilde{f}_{\#}^2(\tilde{E}) = \tilde{E} \cdot \tilde{\mu}_1 \cdot \tilde{\mu}_2$ for some r -legal subpath $\tilde{\mu}_2$ of height r that ends with an edge of H_r . Iterating this produces a nested increasing sequence of paths $\tilde{E} \subset \tilde{f}(\tilde{E}) \subset \tilde{f}_{\#}^2(\tilde{E}) \subset \dots$ whose union is a ray $\tilde{R}$ that converges to some attractor $P \in \text{Fix}_N(\hat{f})$.

The transition matrix for H_r is aperiodic since $f_{\#}(E)$ contains E . Let β be a generic leaf of $\tilde{\Lambda}_r$. Following Definition 3.1.7 of [2], a path of the form $f_{\#}^k(E_i)$ where $k \geq 0$ and E_i is an edge of H_r is called a *tile*. By Lemma 2.9 and by construction, every tile occurs infinitely often in $\tilde{R}$. By Corollary 3.1.11 of [2], each subpath of β is contained in some tile and each tile occurs as a subpath of β . It follows that the accumulation set of P is equal to the weak limit of β which is Λ_r^+ because β is generic. This proves (1).

Assuming now the notation of (2), each finite subpath of $\tilde{R}^{-1}\tilde{R}'$ is contained in a tile. This implies, as in the previous case, that $\tilde{R}^{-1}\tilde{R}'$ is contained in Λ_r and that $\tilde{R}^{-1}\tilde{R}'$ is birecurrent. Lemma 3.1.15 of [2] implies that $\tilde{R}^{-1}\tilde{R}'$ is generic. $\square$

As noted in section 2.4, $\text{Out}(F_n)$ acts on the set of laminations in G . The stabilizer $\text{Stab}(\Lambda)$ of a lamination Λ is the subgroup of $\text{Out}(F_n)$ whose elements leave Λ invariant. We recall Corollary 3.3.1 of [2].

Lemma 2.14. *For each $\Lambda \in \mathcal{L}(\phi)$, there is a homomorphism $PF_{\Lambda} : \text{Stab}(\Lambda) \rightarrow \mathbb{Z}$ such that $\psi \in \text{Ker}(PF_{\Lambda})$ if and only if $\Lambda \notin \mathcal{L}(\psi)$ and $\Lambda \notin \mathcal{L}(\psi^{-1})$.*

We refer to PF_{Λ} as the *expansion factor homomorphism* associated to Λ . The notation is chosen to remind readers that the expansion factor is realized as the logarithm of a Perron Frobenius eigenvalue in a natural way.

2.7 Modifying relative train track maps

To simplify certain arguments in section 3 and as a step toward our ultimate existence theorem (Theorem 4.29), we add properties to the relative train track maps produced in [4]. We can not simply quote results from [2] because, unlike in [2], here we do not allow iteration and we make no assumptions on ϕ .

We need some further notation. The *set of periodic points of f* is denoted $\text{Per}(f)$. A path α is *pre-trivial* if $f_{\#}^k(\alpha)$ is trivial for some $k > 0$. We say that a *nested sequence $\mathcal{C}$ of free factor systems $\mathcal{F}^1 \sqsubset \mathcal{F}^2 \sqsubset \dots \sqsubset \mathcal{F}^m$ is realized* by a relative train track map $f : G \rightarrow G$ and filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ if each $\mathcal{F}^j$ is realized by some $G_{l(j)}$. For any finite graph K , the *core of K* is the subgraph of K consisting of edges that are crossed by some circuit in K . If $K_0 = K$ and K_{i+1} is inductively obtained from K_i by removing all edges with at least one valence one endpoint, then K_i is the core of K for all sufficiently large i .

The transition matrix M_r for an EG stratum H_r of a topological representative $f : G \rightarrow G$ of ϕ has a Perron-Frobenius eigenvalue $\lambda_r > 1$. The set $\{\log(\lambda_r) : H_r \text{ is EG}\}$, listed in non-increasing order, is denoted $\text{PF}(f)$. (In [4] this set is denoted $\Lambda(f)$ but Λ is now usually reserved for laminations.) The set $\{\text{PF}_{\Lambda}(\phi) : \Lambda \in \mathcal{L}(\phi)\}$ of expansion factors for ϕ , listed in non-increasing order, is denoted $\text{EF}(\phi)$. Then $\text{PF}(f) \geq \text{EF}(\phi)$ in the lexicographical order for all f representing ϕ and $\text{PF}(f) = \text{EF}(\phi)$ if $f : G \rightarrow G$ is a relative train track map representing ϕ by Lemma 3.3 of [2].

The number of indivisible Nielsen paths for $f : G \rightarrow G$ with height r is denoted $N_r(f)$.

Remark 2.15. If $f : G \rightarrow G$ satisfies (RTT-*i*) and if $\text{PF}(f) = \text{EF}(\phi)$ then $f : G \rightarrow G$ satisfies (RTT-*iii*) by Lemma 5.9 of [4]. Thus any topological representative that satisfies (RTT-*i*), (RTT-*ii*) and $\text{PF}(f) = \text{EF}(\phi)$ is a relative train track map.

Lemma 2.16. *Suppose that $f : G \rightarrow G$ and $f' : G' \rightarrow G'$ are relative train track maps with EG strata H_r and H'_s respectively and that $p : G \rightarrow G'$ is a homotopy equivalence such that*

- (1) $p(G_r) = G'_s$ and p induces a bijection between the edges of H_r and the edges of H'_s .
- (2) $p_{\#}f_{\#}(\sigma) = f'_{\#}p_{\#}(\sigma)$ for all paths $\sigma \subset G_r$ with endpoints at vertices.

Then $p_{\#}$ induces a bijection between the indivisible periodic Nielsen paths in G with height r and the indivisible periodic Nielsen paths in G' with height s .

Proof. If σ is a periodic Nielsen path for f then $f'_{\#}p_{\#}(\sigma) = p_{\#}f_{\#}(\sigma) = p_{\#}(\sigma)$ which proves that $\sigma' = p_{\#}(\sigma)$ is a periodic Nielsen path for f' . If σ has height r and is indivisible then σ begins and ends in H_r and has exactly one illegal turn in H_r by Lemma 2.11. Our hypotheses imply that the same is true of σ' which implies that σ' is indivisible. For the converse note that for any path $\sigma' \subset G'_s$ that begins with

$p(E_i) \subset H'_s$ and ends with $p(E_j) \subset H'_s$ there is a unique path σ that begins with E_i , ends with E_j and that satisfies $p_{\#}(\sigma) = \sigma'$. By the same reasoning σ is an indivisible Nielsen path if σ' is. $\square$

We next recall the *sliding* operation from [2]. Suppose that H_s is a non-periodic NEG stratum with edges $\{E_1, \dots, E_m\}$ satisfying $f(E_l) = E_{l+1}u_l$ for paths $u_l \subset G_{s-1}$ where $1 \leq l \leq m$ and indices are taken mod m . Choose $1 \leq i \leq m$ and let τ be a path in G_{s-1} from the terminal endpoint v_i of E_i to some vertex w_i . Roughly speaking, we use τ to continuously change the terminal endpoint of E_i from v_i to w_i and to mark the new graph.

More precisely, define a new graph G' from G by replacing E_i with an edge E'_i that has terminal vertex w_i and that has the same initial vertex as E_i . There are homotopy equivalences $p : G \rightarrow G'$ and $p' : G' \rightarrow G$ that are the identity on the common edges of G and G' and that satisfy $p(E_i) = E'_i\bar{\tau}$ and $p'(E'_i) = E_i\tau$. Use p to define the marking on G' and define $f' : G' \rightarrow G'$ on edges by tightening $pf p'$. Complete details can be found in section 5.4 of [2].

Lemma 2.17. *With notation as above.*

- $f' : G' \rightarrow G'$ is a relative train track map.
- $f'|_{G_{s-1}} = f|_{G_{s-1}}$
- If $m = 1$ then $f'(E'_1) = E'_1[\bar{\tau}u_1f(\tau)]$
- If $m \neq 1$ then $f'(E_{i-1}) = E'_i[\bar{\tau}u_{i-1}]$, $f'(E'_i) = E_{i+1}[u_i f(\tau)]$ and $f'(E_j) = E_j u_j$ for $j \neq i-1, i$.
- For each EG stratum H_r , $p_{\#}$ defines a bijection between the set of the indivisible periodic Nielsen paths in G with height r and the indivisible periodic Nielsen paths in G' with height r .

Proof. If $m = 1$ then this is contained in Lemma 5.4.1 of [2]. The argument for $m > 1$ is a straightforward extension of $m = 1$ case and we leave the details to the reader. $\square$

Notation 2.18. Suppose that $u < r$ and that

- (1) H_u is irreducible.
- (2) H_r is EG and each component of G_r is non-contractible.
- (3) for each $u < i < r$, H_i is a zero stratum that is a component of G_{r-1} and each vertex of H_i has valence at least two in G_r .

We say that each H_i is *enveloped by* H_r and write $H_r^z = \cup_{k=u+1}^r H_k$. It is often convenient to treat H_r^z as a single unit.

Theorem 2.19. *For every $\phi \in \text{Out}(F_n)$ there is a relative train track map $f : G \rightarrow G$ and filtration that represents ϕ and satisfies the following properties.*

- (V) *The endpoints of all indivisible periodic Nielsen paths are vertices.*
- (P) *If a stratum $H_m \subset \text{Per}(f)$ is a forest then there exists a filtration element G_j such $\mathcal{F}(G_j) \neq \mathcal{F}(G_l \cup H_m)$ for any G_l . (See also items (1) and (5) of Lemma 2.20.)*
- (Z) *Each zero stratum H_i is enveloped by an EG stratum H_r . Each vertex in H_i is contained in H_r and has link contained in $H_i \cup H_r$.*
- (NEG) *The terminal endpoint of an edge in a non-periodic NEG stratum H_i is periodic and is contained in a filtration element of height less than i that is its own core.*
- (F) *The core of each filtration element is a filtration element.*

Moreover, if $\mathcal{C}$ is a nested sequence of non-trivial ϕ -invariant free factor systems then we may assume that $f : G \rightarrow G$ realizes $\mathcal{C}$.

Before proving Theorem 2.19 we record some useful observations.

Lemma 2.20. *Suppose that $f : G \rightarrow G$ is a relative train track map representing ϕ with filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$.*

- (1) *Suppose that $f : G \rightarrow G$ satisfies (P), that $H_m \subset \text{Per}(f)$ is a forest and that v has valence one in H_m . Then $v \in G_j$ for some $j < m$. If in addition, $f : G \rightarrow G$ satisfies (F) then we may choose G_j to be its own core.*
- (2) *If $f : G \rightarrow G$ satisfies (F) and H_r is EG then G_r is its own core.*
- (3) *If $f : G \rightarrow G$ satisfies (P) and (NEG) then each contractible component of a filtration element is composed of zero strata.*
- (4) *If $f : G \rightarrow G$ satisfies (Z) and (NEG) and if G_k is a filtration element that is its own core, then every vertex in G_k has at least two gates in G_k .*
- (5) *If $f : G \rightarrow G$ satisfies (P) then no component of a filtration element G_m is a forest in $\text{Per}(f)$.*

Proof. Suppose that $f : G \rightarrow G$ satisfies (P) and that $H_m \subset \text{Per}(f)$ is a forest. By (P) there exists j so that $\mathcal{F}(G_j) \neq \mathcal{F}(G_l \cup H_m)$ for any l ; if $f : G \rightarrow G$ also satisfies (F) then we may assume that G_j is its own core. In particular, $\mathcal{F}(G_j \cup H_m) \neq \mathcal{F}(G_j)$. Thus some, and hence every, vertex of valence one in H_m must have valence at least two in $G_j \cup H_m$. This proves (1).

Item (5) follows from the first part of (1).

If H_r is EG then G_r is the smallest filtration element that contains the attracting lamination associated to H_r . This proves (2).

For (3), suppose that C is a contractible component of some G_i . If C contains an edge in an irreducible stratum then it is non-wandering and by (NEG) the lowest stratum H_j in C is either periodic or EG. But H_j can not be periodic by item (1) of this lemma, and cannot be EG by Lemma 2.10. Thus every edge in C is contained in a zero stratum.

The proof of (4) is by induction on k . Suppose that G_k is a filtration element that is its own core and that v is a vertex in G_k . If no illegal turns in G_k are based at v then the number of gates based at v is at least the valence of v in G_k , and so is at least two. We may therefore assume that there is an illegal turn (d_1, d_2) in G_k based at v . By (Z), either both d_1 and d_2 belong to the same EG stratum or one of the d_i 's is the terminal end of a non-fixed NEG edge. Lemma 2.10, (NEG) and the inductive hypothesis imply that in both of these cases v has at least two gates in G_k . $\square$

Proof of Theorem 2.19. By Lemma 2.6.7 of [2], there is a relative train track map $f : G \rightarrow G$ and filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ that represents ϕ and realizes $\mathcal{C}$. (In the statement of Lemma 2.6.7, $\mathcal{C}$ is replaced by a single invariant free factor system $\mathcal{F}$. The more general case that we use is explicitly included in the proof of that lemma.) Lemma 2.12 implies that (V) can be arranged via a finite subdivision. For convenient reference we divide the rest of the proof into steps. Changes are made to $f : G \rightarrow G$ in these steps but we start each step by referring to the current relative train track map as $f : G \rightarrow G$.

If $\mathcal{C}$ has not already been specified, let $\mathcal{C}$ be the nested sequence of free factor systems determined by the $\mathcal{F}(G_r)$'s. For possible future application we will prove the following statement in place of (P).

(P_C) If a stratum $H_m \subset \text{Per}(f)$ is a forest then there exists $\mathcal{F}^j \in \mathcal{C}$ that is not realized by $G_l \cup H_m$ for any G_l .

If $\mathcal{C}$ is determined by the $\mathcal{F}(G_r)$'s then (P) and (P_C) are the same but otherwise the latter is stronger than the former.

Remark 2.21. For reference in the proof of Theorem 4.29 we record the following property of the remainder of our construction. If $f : G \rightarrow G$ is the relative train track map as it is now and $g : G' \rightarrow G'$ is the ultimate modified relative train track map produced by the six steps listed below, then there is a bijection $H_r \rightarrow H'_s$ between the EG strata of $f : G \rightarrow G$ and the EG strata of $g : G' \rightarrow G'$ such that

- (a) H_r and H'_s have the same number of edges.
- (b) $N_r(f) = N_s(g)$.

The five moves used in our construction are valence two homotopies away from EG strata, sliding (which is defined following Lemma 2.16), reordering of strata, tree replacements (see step 2) and collapsing forests in NEG and zero strata. Item (a) will be obvious as will (b) for the valence two homotopies and the reordering of strata. For the remaining three moves (b) will follow from Lemmas 2.17 and 2.16.

(Step 1 : A weak form of (NEG)) Suppose that H_s is a non-periodic NEG stratum with edges $\{E_1, \dots, E_m\}$ satisfying $f(E_i) = E_{i+1}u_i$ for paths $u_i \subset G_{s-1}$ where indices are taken mod m . Our goal in this step is to arrange

- (1) The endpoints of an NEG edge are periodic.

Care is taken that no vertices with valence one in G are created.

We first arrange that the terminal endpoint v_1 of E_1 is either periodic or has valence at least three in G . If this is not already the case, let E be the unique oriented edge of G , other than E_1 , whose initial endpoint is v_1 . Since H_s is not a periodic stratum, $E \subset G_{s-1}$. Lemma 2.10 implies that E does not belong to an EG stratum. Perform a valence two homotopy as defined on page 13 of [4]. There are two steps to this operation. The first is to modify f so that no vertex is mapped to v_1 . The second is to amalgamate E_1E into a single edge named E_1 by removing v_1 from the list of vertices. If σ is a path, neither of whose endpoints are mapped to v_1 by the original map, then occurrences of E_1 and E in the original edge path for $f_{\#}(\sigma)$ always occur as E_1E or its inverse; the new edge path for $f_{\#}(\sigma)$ is obtained from the original by removing all occurrences of E . Since v_1 is not incident to any EG stratum, this applies to any path σ with endpoints in EG strata. It follows that the new map (still called $f : G \rightarrow G$) is still a relative train track map, satisfies (V) and realizes $\mathcal{C}$. It also follows that this operation does not change the number of edges in any EG stratum and does not change the number of indivisible Nielsen paths with height corresponding to any EG stratum. After finitely many valence two homotopies we may assume that the terminal endpoint v_1 of E_1 is either periodic or has valence at least three in G .

The component C_1 of G_{s-1} that contains v_1 does not wander and so contains a periodic vertex w_1 . Choose a path $\tau \subset G_{s-1}$ from v_1 to w_1 in C and slide to change the terminal endpoint of E_1 to w_1 . Because of our previous move, v_1 still has valence at least two in G .

After repeating these operations finitely many times we have arranged (1) and not created any valence one vertices.

(Step 2 : A weak form of (Z)) In this step we prove most of (Z). The one missing item is that we only show that the components of G_r are non-wandering instead of showing that they are non-contractible.

The union of the non-wandering components of a filtration element G_i is f -invariant. Thus the strata, if any, that are contained in wandering components of G_i can be moved up the filtration to be above the strata contained in the non-wandering components of G_i . After finitely many such changes and after amalgamating zero strata if necessary, we may assume that

- (2) If G_i has wandering components then H_i is a wandering component of G_i .

Suppose that K is a component of the union of all zero strata, that H_i is the highest stratum that contains an edge in K and that H_u is the highest irreducible

stratum below H_i . We prove that $K \cap G_u = \emptyset$ by assuming that $K \cap G_u \neq \emptyset$ and arguing to a contradiction. Since each component of G_u is non-wandering and since some iterate of f maps K into G_u , there is a unique component C of G_u that intersects K . If each vertex $v \in K$ has valence at least two in $C \cup K$ then each edge of K is contained in a path in K with endpoints in C , and so by the connectivity of C , is contained in a circuit in $K \cup C$. This contradicts the fact that some iterate of f maps $K \cup C$ into C , and we conclude that some vertex v of K has valence one in $K \cup C$. In particular, v is non-periodic and so is not the endpoint of an NEG edge by (1) and is not the endpoint of an EG edge in a stratum above G_i by Remark 2.8. By construction, v is not the endpoint of an edge in a zero stratum above G_i . But then v has valence one in G . This contradiction proves that $K \cap G_u = \emptyset$. After reorganizing the edges in zero strata, we may assume that $H_i = K$. Note in particular that no vertex in H_i is periodic.

Let H_r be the first irreducible stratum above H_i . The component of G_r that contains H_i is non-wandering by (2) and so must intersect H_r . Since no vertex of H_i is periodic, (1) implies that H_r is EG. Moreover, the argument used in the previous paragraph proves that the link in G of each vertex in H_i is contained in $H_i \cup H_r \subset G_r$ and that H_i is contained in the core of G_r .

We arrange that each vertex in H_i is contained in H_r by the following *tree replacement* move. Replace H_i with a tree H'_i whose vertex set is exactly $H_i \cap H_r$. Do this for each zero stratum H_i and call the resulting graph G' . We view the union X of the irreducible strata as a subgraph of both G and G' . The set of vertices in X is f -invariant by (RTT-*i*) and (1). There is a homotopy equivalence $p' : G' \rightarrow G$ that is the identity on X and that sends an edge E' of H'_i to the unique path in H_i connecting the endpoints of E' . The homotopy inverse $p : G \rightarrow G'$ is the identity on X and sends H_i to H'_i ; the exact definition depends on choosing $p(w) \in H'_i$ for those vertices $w \in H_i$, if any, that are not contained in X . Note that $p_\#$ defines a bijection, with inverse $p'_\#$, between paths in G with endpoints at vertices in X and paths in G' with endpoints at vertices in X . The homotopy equivalence $f' : G' \rightarrow G'$ defined on edges by $f'(E') = (pfp')_\#(E')$ is independent of the choices made in defining p and represents ϕ . To verify (RTT-*ii*) for f' it suffices to show that if $\alpha' \subset H'_i$ is non-trivial then $f'_\#(\alpha') = p_\#f_\#(p'_\#(\alpha'))$ is non-trivial. This follows from (RTT-*ii*) for f and the fact that both $p_\#$ and $p'_\#$ preserve non-triviality for paths with endpoints in X . It is easy to see that (RTT-*i*) for f implies (RTT-*i*) for f' and that $\text{PF}(f) = \text{PF}(f')$. Remark 2.8 implies that $f' : G' \rightarrow G'$ is a relative train track map. None of the moves in this step change the free factor systems represented by filtration elements so $f' : G' \rightarrow G'$ still realizes $\mathcal{C}$.

(Step 3 : Property (P_C)) If (P_C) fails then there is a stratum $H_m \subset \text{Per}(f)$ that is a forest with the property that for each $\mathcal{F}^j \in \mathcal{C}$ there is a filtration element $G_{l(j)}$ such that $G_{l(j)} \cup H_m$ realizes $\mathcal{F}^j$. We will construct a new relative train track map $f' : G' \rightarrow G'$ with one fewer NEG stratum that still realizes $\mathcal{C}$ and satisfies (1) and

the weak form of (Z). After repeating this finitely many times we will have achieved (Pc).

Let Y be the set of edges in $G \setminus H_m$ that are mapped entirely into H_m by some iterate of f . Then each edge of Y is contained in a zero stratum and $H_m \cup Y$ is a forest that is mapped into itself by f and into H_m by some iterate of f . We next arrange that

- (*) if α is a path in a zero stratum with endpoints at vertices and if α is not contained in Y then $f_{\#}(\alpha)$ is not contained in $H_m \cup Y$.

Suppose to the contrary that $\alpha \subset H_k$ violates (*). Choose an edge $E_i \subset H_k$ that is crossed by α and is not contained in Y . Perform a tree replacement move on H_k as in step 2, replacing E_i by an edge connecting the endpoints of α . The new edge is mapped entirely into H_m by some iterate of f and we add it to Y . After finitely many such moves (*) is satisfied.

Let G' be the marked graph obtained by collapsing each component of $H_m \cup Y$ to a point and let $p : G \rightarrow G'$ be the corresponding quotient map. Identify the edges of G' with those of $G \setminus (H_m \cup Y)$ and define $f' : G' \rightarrow G'$ by $f'(E) = [pf(E)]$. As an edge path, $f'(E)$ is obtained from $f(E)$ by removing all occurrences of edges in $H_m \cup Y$. It follows that the strata H_r and $p(H_r)$ (if the latter is non-empty) have the same type (zero, EG, NEG), that $f' : G' \rightarrow G'$ has one fewer NEG stratum than $f : G \rightarrow G$, that $f' : G' \rightarrow G'$ satisfies (1) and the weak form of (Z), that $\text{PF}(f') = \text{PF}(f)$ and that $f' : G' \rightarrow G'$ satisfies (RTT-*i*). Lemma 5.9 of [4] implies that $f' : G' \rightarrow G'$ satisfies (RTT-*iii*).

To verify (RTT-*ii*), suppose that $H'_s = p(H_r)$ is EG and that $\alpha' \subset G'_{s-1}$ is a connecting path for H'_s . If α' is contained in a zero stratum then $f'_{\#}(\alpha')$ is non-trivial by (*). We may therefore assume that the component C' of G'_{s-1} that contains α' is not a zero stratum and hence is non-wandering. To prove that $f'_{\#}(\alpha')$ is non-trivial, it suffices, by Remark 2.8, to show that each $v' \in H'_s \cap C'$ is a periodic point.

Since v' is incident to an edge in H'_s , there is a vertex $v \in H_r$ such that $p(v) = v'$. If v is periodic, we are done. We may therefore assume that $v \notin H_m$. If $v \in Y$ then the component of G_{r-1} that contains v is a zero stratum by the weak form of (Z) contradicting the assumption that v' is contained in a non-wandering component of G'_{s-1} . It follows that $v = p^{-1}(v')$. By the same reasoning, v is contained in a non-wandering component of G_{r-1} and so is periodic by Remark 2.8. Thus v' is periodic and we have verified (RTT-*ii*) for f' . This completes the proof that $f' : G' \rightarrow G'$ is a relative train track map.

There exists $k > 0$ so that each non-contractible component of $G_{l(j)} \cup H_m$ is f^k -invariant and so that f^k induces a rank preserving bijection between the non-contractible components of $G_{l(j)} \cup H_m \cup Y$ and the non-contractible components of $G_{l(j)} \cup H_m$. Thus $G_{l(j)} \cup H_m \cup Y$ and hence $p(G_{l(j)})$ realizes $\mathcal{F}^j$, proving that $f' : G' \rightarrow G'$ and its filtration realize $\mathcal{C}$.

(Step 4 : Property (Z)) The lowest stratum in a non-wandering component C of a filtration element is either EG or periodic. Lemma 2.10 and item (5) of Lemma 2.20 imply that C is not contractible. (Z) therefore follows from the weak form of (Z).

(Step 5 : Property (NEG)) Suppose that H_s is a non-periodic NEG stratum with edges $\{E_1, \dots, E_m\}$ satisfying $f(E_i) = E_{i+1}u_i$ for paths $u_i \subset G_{s-1}$ where indices are taken mod m . The component C_i of G_{s-1} that contains the terminal endpoint v_i of E_i does not wander. The lowest stratum H_t in C_i is either EG or periodic. In the former case, every vertex in H_t has at least two gates in H_t by Lemma 2.10 and so H_t is its own core. In the latter case, the same result follows from (P_C) and item (1) of Lemma 2.20.

Choose a path $\tau \subset G_{s-1}$ from v_i to a periodic vertex w_i in H_t and slide to change the terminal endpoint of E_i to w_i . After performing this sliding operation finitely many times, working up through the filtration, (NEG) is satisfied. The resulting homotopy equivalence is a relative train track map by Lemma 2.17, still realizes $\mathcal{C}$ and still satisfies (Z).

Sliding may have introduced valence one vertices to G . But no such vertex is the image of a vertex with valence greater than one by (NEG), (Z) and Lemma 2.10. We may therefore remove all vertices of valence one and the edges that are incident to them. After repeating this finitely many times G has no valence one vertices.

If (P_C) is no longer satisfied then return to step 3. Since this reduces the number of NEG stratum the process stops.

(Step 6 : Property (F)) If H_l is a zero stratum then G_l and G_{l-1} realize the same free factor system. We may therefore assume that G_l is not a zero stratum and hence that every component of G_l is non-contractible. If w is a valence one vertex of G_l then by item (1) of Lemma 2.20, (NEG) and Lemma 2.10, w must be the initial endpoint of a non-periodic NEG edge in some H_k with $k \leq l$ and no vertex with valence at least two in G_l maps to w . The initial endpoint of each edge in H_k has valence one in G_l and $G_l \setminus H_k$ is f -invariant. We may therefore reorder the strata to move H_k above $G_l \setminus H_k$. After finitely many such moves, G_l is a core graph. Working our way up the filtration we arrange that (F) is satisfied. $\square$

We conclude this section by recalling an operation from page 46 of [4] and Definitions 5.3.2 of [2].

Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ that satisfies item (Z) of Theorem 2.19 and that ρ is an indivisible Nielsen path of height r . Decompose $\rho = \alpha\beta$ into a concatenation of maximal r -legal subpaths as in Lemma 2.11 and let $E_1 \subset H_r$ and $E_2 \subset H_r$ be the initial edges of $\bar{\alpha}$ and β respectively. If one of the edge paths $f(E_i)$, $i=1$ or 2 , is an initial subpath of the other then we say that *the fold at the illegal turn of ρ is a full fold*; otherwise it is a *partial fold*. There are two kinds of full folds. If $f(E_1) \neq f(E_2)$ then the full fold is *proper*; otherwise it is *improper*.

Suppose that the fold at the illegal turn of ρ is proper, say that $f(E_1)$ is a proper

initial subpath of $f(E_2)$. Write $\bar{\alpha} = E_1 b E_3 \dots$ where b is a (possibly trivial) subpath of G_{r-1} and E_3 is an edge in H_r . Since both the initial edge of $f(E_3)$ and the first edge of $f(\beta)$ that is not cancelled when $f(\alpha)f(\beta)$ is tightened to $\alpha\beta$ belong to H_r , there is a decomposition $E_2 = E_2'' E_2'$ into subpaths such that $f(E_2'') = f_{\#}(E_1 b)$ and such that the first edge in $f(E_2')$ is contained in H_r . Subdivide E_2 into two edges E_2'' and E_2' and then identify E_2'' with $E_1 b$ to form a new graph G' . The quotient map $F : G \rightarrow G'$ is called the *extended fold determined by ρ_r* .

We think of $G \setminus E_2$ as a subgraph of G' on which F is the identity. By construction $F(E_2) = E_1 b E_2'$. The filtration on G' is defined by $H'_i = H_i$ for $i \neq r$ and $H'_r = (H_r \setminus E_2) \cup E_2'$. There is a map $g : G' \rightarrow G$ such that $gF = f$. We refer to $g : G' \rightarrow G$ as *map induced by the extended fold*.

The following lemma states that the map $f' : G' \rightarrow G'$ obtained from $Fg : G' \rightarrow G'$ by tightening the images of edges is a relative train track map that satisfies item (Z) of Theorem 2.19. We say that $f' : G' \rightarrow G'$ is obtained from $f : G \rightarrow G$ by *folding ρ_r* and that $\rho'_r = F_{\#}(\rho_r)$ is *the indivisible Nielsen path determined by ρ_r* . If the illegal turn of ρ'_r is proper then this process can be repeated. This is referred to as *iteratively folding ρ* .

Lemma 2.22. *Assuming notation as above, $f' : G' \rightarrow G'$ is a relative train track map that satisfies item (Z) of Theorem 2.19.*

Proof. By construction, $f'|G_{r-1} = f|G_{r-1}$. If E is an edge in H_r then $f(E)$ does not cross the illegal turn in ρ_r . If $E \neq E_2$ then $f'(E)$ is obtained from $f(E)$ by replacing each occurrence of E_2 with $E_1 b E_2''$. Similarly, $f'(E_2')$ is obtained from $f(E_2')$ by replacing each occurrence of E_2 with $E_1 b E_2''$. It follows that H'_r satisfies (RTT-i) - (RTT-iii).

If H_k is a zero stratum above H_r then each edge E_k in H_k is a connecting path for some EG stratum H_s above H_k by item (Z) of Theorem 2.19. Thus $f_{\#}^i(E_k)$ is non-trivial for all $i \geq 0$. Since F does not identify points that are not identified by f , and since $F|E_k$ is the identity, $(Fg)_{\#}(E_k) = (Ff)_{\#}(E_k)$ is non-trivial. This shows that no edges are collapsed when Fg is tightened to f' . The same argument shows that if $\sigma \subset H_k$ is any path with endpoints at vertices then $f'_{\#}(\sigma)$ is non-trivial.

If H_l is NEG then H'_l is NEG.

Suppose that E_m is an edge in an EG stratum H_m above H_r and that $f(E_m) = \mu_1 \nu_1 \mu_2 \dots \nu_l \mu_{l+1}$ is the decomposition into subpaths $\mu_j \subset H_m$ and subpaths $\nu_j \subset G_{m-1}$. Then $f'(E_m) = (Fg)_{\#}(E_m) = (Ff)_{\#}(E_m) = \mu_1 \nu'_1 \mu_2 \dots \nu'_l \mu_{l+1}$ where $\nu'_j = F_{\#}(\nu_j)$ is non-trivial because $f_{\#}(\nu_j)$ is non-trivial. This proves that H'_m satisfies (RTT-i) and (RTT-iii).

To verify (RTT-ii) for H_m suppose that σ' is a connecting path for H'_m . If σ' is contained in a zero stratum H'_k then it is disjoint from G_r and so is identified with a connecting path $\sigma \subset H_k$. By our previous argument, $f'_{\#}(\sigma')$ is non-trivial. If σ' is contained in non-contractible component of G'_{m-1} then there is a connecting path σ for H_m in a non-contractible component of G_{m-1} such that $F_{\#}(\sigma) = \sigma'$. The

endpoints of σ are periodic for f by Remark 2.8. It follows that the endpoints of σ' are periodic for f' and another application of Remark 2.8 proves that H'_m satisfies (RTT-ii). This completes the proof that $f' : G' \rightarrow G'$ is a relative train track map.

Item (Z) of Theorem 2.19 for f' therefore follows from item (Z) of Theorem 2.19 for f . $\square$

3 Forward Rotationless Outer Automorphisms

To avoid issues raised by finite order phenomenon, one often replaces $\phi \in \text{Out}(F_n)$ with an iterate ϕ^k . In this section we explain how this can be done canonically by exhibiting the natural class of outer automorphisms that require no iteration. We also define principal automorphisms in the context of $\text{Out}(F_n)$. These automorphisms play a central role in both the definition of forward rotationless outer automorphisms (Definition 3.13) and in the formulation of the Recognition Theorem (Theorem 5.3).

In section 3.1, we recall how principal automorphisms occur in the context of the mapping class group. Examples and definitions for $\text{Out}(F_n)$ are given in section 3.2. An equivalent definition is then given in terms of relative train track maps and the Nielsen classes of their fixed points. Finally, in section 3.5 we record some properties of forward rotationless outer automorphisms that justify their name; for example, we show that a ϕ -periodic free factor is ϕ -invariant.

3.1 The Nielsen Approach to the Mapping Class Group

To provide historical context and motivation for our techniques and results, we briefly recall Nielsen's point of view on the mapping class group. Further details and proofs can be found, for example, in [13].

Let S be a closed orientable surface of negative Euler characteristic and let $h : S \rightarrow S$ be a homeomorphism representing an element $\mu \in \text{MCG}(S)$. A choice of complete hyperbolic structure on S identifies the universal cover $\tilde{S}$ of S with the hyperbolic plane $\mathbb{H}$. Using the Poincaré disk model for $\mathbb{H}$, there is an induced compactification of $\tilde{S}$ by adding a topological circle S_∞ .

To avoid cumbersome superscripts we use g to denote a positive iterate $g := h^k$ of h . Any lift $\tilde{g} : \tilde{S} \rightarrow \tilde{S}$ of g extends to a homeomorphism of the compactification. The restriction of this extension to S_∞ , denoted $\hat{g} : S_\infty \rightarrow S_\infty$, depends only on k , the isotopy class of h and the choice of lift. More precisely, h induces an outer automorphism of $\pi_1(S)$ and $\hat{g} = \widehat{\Phi_{\tilde{g}}}$ where $\Phi_{\tilde{g}}$ is the automorphism of $\pi_1(S)$ corresponding to $\tilde{g}$ and $\widehat{\Phi_{\tilde{g}}} : S_\infty \rightarrow S_\infty$ is the homeomorphism determined by the identification of S_∞ with $\partial\pi_1(S)$ [11].

Denote the set of non-repelling fixed points of $\hat{g}$ by $\text{Fix}_N(\hat{g})$. If $\text{Fix}_N(\hat{g})$ contains at least three non-repelling points then we say that $\tilde{g}$ is a *principal lift* of g and that $\Phi_{\tilde{g}}$ is a principal automorphism representing μ^k . The sets $\text{Fix}_N(\hat{g})$ determined by the principal lifts of iterates of h are central to Nielsen's investigations.

The mapping class μ determined by h is rotationless as defined in the Section 1 if and only for all k , each principal lift $\tilde{g}$ of $g = h^k$ has the form $\tilde{h}^k$ where $\tilde{h}$ is a principal lift of h and where $\text{Fix}_N(\hat{g}) = \text{Fix}_N(\hat{h})$. Thus from the point of view of principal lifts and their Fix_N sets, nothing changes if μ is replaced by an iterate. For the remainder of this discussion we assume that μ is rotationless and that $k = 1$.

The intersection $\tilde{\Delta}(\tilde{g})$ of the convex hull of $\text{Fix}_N(\hat{g})$ with $\tilde{S} = \mathbb{H}$ is called the *principal region* for $\tilde{g}$ and its image in S is denoted $\Delta(\tilde{g})$. Thus $\tilde{g}$ is principal if and only if $\Delta(\tilde{g})$ has non-empty interior.

Assume that $\tilde{g}$ is principal. If no point in $\text{Fix}_N(\hat{g})$ is isolated then $\Delta(\tilde{g})$ is a compact subsurface and there is a homeomorphism $f : S \rightarrow S$ representing μ whose restriction to $\Delta(\tilde{g})$ is the identity. If $\text{Fix}_N(\hat{g})$ is finite, or more generally, is finite up to the action of a single covering translation that commutes with $\tilde{g}$, then the interior of $\Delta(\tilde{g})$ is a component of the complement in S of one of the pseudo-Anosov laminations Λ associated to μ . The boundary of $\Delta(\tilde{g})$ is a finite union of leaves of Λ and perhaps one reducing curve. These are the only cases that occur if there is non-trivial twisting along each reducing curve in the Thurston normal form for μ . In the general case, $\Delta(\tilde{g})$ is a finite union of the two types.

A proof of (most of) the Thurston classification theorem from this point of view is contained in [13] and [16].

3.2 Principal Automorphisms

Suppose that $f : G \rightarrow G$ is a relative train track map representing $\phi \in \text{Out}(F_n)$. Recall from section 2.3 that there is a bijection between lifts $\tilde{f} : \Gamma \rightarrow \Gamma$ to the universal cover and automorphisms $\Phi \in \text{Aut}(F_n)$ representing ϕ . A fixed point

Definition 3.1. For $\Phi \in \text{Aut}(F_n)$ representing ϕ , let $\text{Fix}_N(\hat{\Phi}) \subset \text{Fix}(\hat{\Phi})$ be the set of non-repelling fixed points of $\hat{\Phi}$. We say that Φ is a *principal automorphism* and write $\Phi \in \text{P}(\phi)$ if either of the following hold.

- $\text{Fix}_N(\hat{\Phi})$ contains at least three points.
- $\text{Fix}_N(\hat{\Phi})$ is a two point set that is neither the set of endpoints of an axis A_c nor the set of endpoints of a lift $\tilde{\lambda}$ of a generic leaf of an element of $\mathcal{L}(\phi)$.

The corresponding lift $\tilde{f} : \Gamma \rightarrow \Gamma$ is a *principal lift*.

Remark 3.2. For all $\phi \in \text{Out}(F_n)$ there exists, by Lemma 5.2 of [3] or Proposition I.5 of [14], $k \geq 1$ such that $\text{P}(\phi^k) \neq \emptyset$. Moreover, if the conjugacy class of $a \in F_n$ is invariant under ϕ^k , then one may choose $\Phi \in \text{P}(\phi^k)$ to fix a .

Remark 3.3. If $\text{Fix}(\Phi)$ has rank at least two then Φ is a principal automorphism by Lemma 2.3.

Remark 3.4. If Φ_1 and Φ_2 are distinct representatives of ϕ then $\text{Fix}_N(\hat{\Phi}_1) \cap \text{Fix}_N(\hat{\Phi}_2)$ is contained in $\text{Fix}(\hat{\Phi}_1^{-1}\hat{\Phi}_2) = \{T_c^\pm\}$ for some non-trivial covering translation T_c . It follows that if Φ_1 and Φ_2 are principal then $\text{Fix}_N(\hat{\Phi}_1) \neq \text{Fix}_N(\hat{\Phi}_2)$.

Remark 3.5. The second item in our definition of principal automorphism does not occur in the context of mapping class groups. It arises in $\text{Out}(F_n)$ to account for nonlinear NEG strata as illustrated by Example 3.10 below.

Remark 3.6. Each $\Lambda \in \mathcal{L}(\phi)$ has infinitely many generic leaves that are invariant by an iterate of $\phi_\#$. If $\{P, Q\}$ is the endpoint set of a lift of such a leaf then (Lemma 4.39) there exists Φ representing an iterate of ϕ such that P and Q are attracting fixed points for $\hat{\Phi}$. Remark 3.9 and Lemma 2.3 imply that for all but finitely many such leaves, $\text{Fix}_N(\hat{\Phi}) = \{P, Q\}$ and Φ is not principal.

Remark 3.7. If Φ has positive index in the sense of [12], then Φ is a principal automorphism. The converse fails for the principal automorphism Φ_2 of Example 3.10.

We say that $x, y \in \text{Fix}(f)$ are *Nielsen equivalent* or belong to the same *Nielsen class* if they are the endpoints of a Nielsen path for f . Each Nielsen class is an open subset of $\text{Fix}(f)$ because every sufficiently short path with endpoints in $\text{Fix}(f)$ is a Nielsen path. In particular, there are only finitely many Nielsen classes.

If $\tilde{f} : \Gamma \rightarrow \Gamma$ is a lift of $f : G \rightarrow G$, then any path $\tilde{\alpha} \subset \Gamma$ with endpoints in $\text{Fix}(\tilde{f})$ projects to a Nielsen path $\alpha \subset G$ for f . Conversely, if α is a Nielsen path for f and $\tilde{f}$ fixes one endpoint of a lift $\tilde{\alpha}$ of α then $\tilde{f}$ also fixes the other endpoint of $\tilde{\alpha}$. Thus $\text{Fix}(\tilde{f})$ is either empty or projects onto a single Nielsen class in $\text{Fix}(f)$.

A pair of automorphisms Φ_1 and Φ_2 are *equivalent* if there exists $c \in F_n$ such that $\Phi_2 = i_c \Phi_1 i_c^{-1}$. Translating this into the language of lifts, $\tilde{f}_1$ is equivalent to $\tilde{f}_2$ if $\tilde{f}_2 = T_c \tilde{f}_1 T_c^{-1}$. This equivalence relation is called *isogredience*.

Lemma 3.8. Suppose that $f : G \rightarrow G$ represents $\phi \in \text{Out}(F_n)$ and that $\tilde{f}_1$ and $\tilde{f}_2$ are lifts of f with non-empty fixed point sets. Then $\tilde{f}_1$ and $\tilde{f}_2$ belong to the same isogredience class if and only if $\text{Fix}(\tilde{f}_1)$ and $\text{Fix}(\tilde{f}_2)$ project to the same Nielsen class in $\text{Fix}(f)$.

Proof. If $\tilde{f}_2 = T_c \tilde{f}_1 T_c^{-1}$ then $\text{Fix}(\tilde{f}_2) = T_c \text{Fix}(\tilde{f}_1)$ and $\text{Fix}(\tilde{f}_2)$ and $\text{Fix}(\tilde{f}_1)$ project to the same Nielsen class in $\text{Fix}(f)$. Conversely, if $\text{Fix}(\tilde{f}_2)$ and $\text{Fix}(\tilde{f}_1)$ have the same non-trivial projection then there exists $\tilde{x} \in \text{Fix}(\tilde{f}_2)$ and a covering translation T_c such that $T_c(\tilde{x}) \in \text{Fix}(\tilde{f}_1)$ which implies that $\tilde{f}_2$ and $T_c \tilde{f}_1 T_c^{-1}$ agree on a point and hence are equal. $\square$

Remark 3.9. We show below (Corollary 3.17) that principal lifts have non-trivial fixed point sets in Γ . Since there are only finitely many Nielsen classes in $\text{Fix}(f)$, it follows that there are only finitely many isogredience classes of principal lifts for ϕ .

In the following examples, G is the rose R_3 with basepoint v at the unique vertex. We use A, B and C to denote both the oriented edges of G and the corresponding

generators of F_3 . Our examples are all positive automorphisms Φ , meaning that they are defined by $A \mapsto w_A$, $B \mapsto w_B$ and $C \mapsto w_C$ where w_A, w_B and w_C are words in the letters A, B and C (and not the inverses $\bar{A}, \bar{B}$ and $\bar{C}$). These words also define a homotopy equivalence $f : G \rightarrow G$. Since w_A, w_B and w_C use only A, B and C , and not $\bar{A}, \bar{B}$ and $\bar{C}$, the homotopy equivalence is a relative train track map for ϕ .

The universal cover of $\tilde{G}$ is denoted Γ and we assume that a basepoint $\tilde{v}$ has been chosen. Some statements in the examples are left for the reader to verify or follow from results we establish later in this section; none of these statements are ever quoted.

Example 3.10. Let $\Phi_1 \in \text{Aut}(F_3)$ be determined by $w_A = A, w_B = BA$ and $w_C = BCB^2$. Then $\text{Fix}(\Phi_1) = \langle A, BAB \rangle$ and $\text{Fix}(\hat{\Phi}_1) = \partial(\text{Fix}(\Phi_1))$. The lift $\tilde{f}_1$ that fixes $\tilde{v}$ is the principal lift corresponding to Φ_1 .

The unique fixed point x of f in the interior of C is not Nielsen equivalent to v . Let $\tilde{C}$ be the lift of C whose initial endpoint is $\tilde{v}$ and let $\tilde{f}_2$ be the lift that fixes the unique lift of $\tilde{x}$ of x in $\tilde{C}$. Then $\tilde{f}_2$ is principal and $\text{Fix}_N(\tilde{f}_2)$ is a pair of attractors which bound the line that is the union of the increasing sequence $\tilde{C} \subset (\tilde{f}_2)_\#(\tilde{C}) \subset (\tilde{f}_2)_\#^2(\tilde{C}) \subset \dots$. If Φ_2 is the principal automorphism corresponding to $\tilde{f}_2$ then $\Phi_2 = i_B^{-1}\Phi_1$.

Example 3.11. Let $\Phi_1 \in \text{Aut}(F_3)$ be determined by $w_A = A, w_B = BA$ and $w_C = CB^2$. Then $\text{Fix}(\Phi) = \langle A, BAB \rangle$ and $\text{Fix}(\hat{\Phi})$ is the union of $\partial(\text{Fix}(\Phi))$ with the $\text{Fix}(\Phi)$ -orbit of a single attractor P . The lift $\tilde{f}$ that fixes $\tilde{v}$ is the principal lift corresponding to Φ and P is the endpoint of the ray that is the union of the increasing sequence $\tilde{C} \subset (\tilde{f}_2)_\#(\tilde{C}) \subset (\tilde{f}_2)_\#^2(\tilde{C}) \subset \dots$.

Example 3.12. Let $\Phi \in \text{Aut}(F_3)$ be determined by $w_A = ACBA, w_B = BA$ and $w_C = CBA$, let $\tilde{f}$ be the lift that fixes $\tilde{v}$ and let $\tilde{A}, \tilde{B}, \tilde{C}$ and $\tilde{A}^{-1}$ be the lifts of the oriented edges A, B, C and $\bar{A}$ with $\tilde{v}$ as initial vertex. The directions determined by the initial edges of $\tilde{A}, \tilde{B}, \tilde{C}$ and $\tilde{A}^{-1}$ are fixed by $D\tilde{f}$. Lemma 2.13 produces attractors P_A, P_B, P_C and $P_{\bar{A}}$ in $\text{Fix}_N(\hat{f})$ such that lines connecting $P_{\bar{A}}$ to the other three points are generic leaves of an attracting lamination. Lemma 3.21 implies that $\text{Fix}_N(\hat{f}) = \{P_A, P_B, P_C, P_{\bar{A}}\}$.

We now come to the second main definition of this section. Note that if Φ is a principal lift of ϕ then Φ^k is a principal lift for ϕ^k and $\text{Fix}_N(\hat{\Phi}) \subset \text{Fix}_N(\hat{\Phi}^k)$ for all $k \geq 1$. The set of non-repelling periodic points in $\text{Per}(\hat{\Phi})$ is denoted $\text{Per}_N(\hat{\Phi})$. By iterating ϕ we might pick up more principal lifts and principal lifts might pick up more non-repelling fixed points. If this doesn't happen, then we say that ϕ is *forward rotationless*. Here is the precise definition.

Definition 3.13. An outer automorphism ϕ is *forward rotationless* if $\text{Fix}_N(\hat{\Phi}) = \text{Per}_N(\hat{\Phi})$ for all $\Phi \in \text{P}(\phi)$ and if for each $k \geq 1$, $\Phi \mapsto \Phi^k$ defines a bijection (see Remark 3.14) between $\text{P}(\phi)$ and $\text{P}(\phi^k)$. Our standing assumption is that $n \geq 2$. For notational convenience we also say that the identity element of $\text{Out}(F_1)$ is forward rotationless.

Remark 3.14. By Remark 3.4 there is no loss in replacing the assumption that $\Phi \mapsto \Phi^k$ defines a bijection with the a priori weaker assumption that $\Phi \mapsto \Phi^k$ defines a surjection.

3.3 Rotationless Relative Train Track Maps and Principal Periodic Points

We now want to characterize those relative train track maps $f : G \rightarrow G$ that represent forward rotationless $\phi \in \text{Out}(F_n)$ and to determine which lifts of such f are principal. We precede our main definitions by showing that principal lifts have fixed points.

Suppose that $f : G \rightarrow G$ represents ϕ and that $\tilde{f} : \Gamma \rightarrow \Gamma$ is a lift of f . We say that $\tilde{z} \in \Gamma$ *moves toward* $P \in \text{Fix}(\hat{f})$ under the action of $\tilde{f}$ if the ray from $\tilde{f}(\tilde{z})$ to P does not contain $\tilde{z}$. Similarly, we say that $\tilde{f}$ *moves* $\tilde{y}_1$ and $\tilde{y}_2$ *away from each other* if the path in Γ connecting $\tilde{f}(\tilde{y}_1)$ to $\tilde{f}(\tilde{y}_2)$ contains $\tilde{y}_1$ and $\tilde{y}_2$ and if $\tilde{f}(\tilde{y}_1) < \tilde{y}_1 < \tilde{y}_2 < \tilde{f}(\tilde{y}_2)$ in the order induced by the orientation on that path.

The following lemma relates the action of $\hat{f}$ to the action of $\tilde{f}$ and gives a criterion for elements of $\text{Fix}(\hat{f})$ to be contained in $\text{Fix}_N(\hat{f})$. Recall that ∂F_n is identified with the set of ends of Γ . It therefore makes sense to say that points in Γ are close to $P \in \partial F_n$ or that P is the limit of points in Γ .

Lemma 3.15. *Suppose that $P \in \text{Fix}(\hat{f})$ and that there does not exist $c \in F_n$ such that $\text{Fix}(\hat{f}) = \{T_c^\pm\}$.*

- (1) *If P is an attractor for the action of $\hat{f}$ on $\partial\Gamma$ then $\tilde{z}$ moves toward P under the action of $\tilde{f}$ for all $\tilde{z} \in \Gamma$ that are sufficiently close to P .*
- (2) *If P is an endpoint of an axis A_c or if P is the limit of points in Γ that are either fixed by $\tilde{f}$ or that move toward P under the action of $\tilde{f}$, then $P \in \text{Fix}_N(\hat{f})$.*

Proof. The lemma is an immediate consequence of Proposition 1.1 of [12] if P is not the endpoint $T_c^\pm$ of an axis A_c . If P is T_c^+ or T_c^- , then $\text{Fix}(\hat{f})$ contains $\{T_c^\pm\}$ and at least one other point. Lemma 2.3 implies that P is not isolated in $\text{Fix}(\hat{f})$ and is therefore neither an attractor nor a repeller for the action of $\hat{f}$. $\square$

The next lemma is based on Lemma 2.1 of [4].

Lemma 3.16. *If $\tilde{f}$ moves $\tilde{y}_1$ and $\tilde{y}_2$ away from each other, then $\tilde{f}$ fixes a point in the interval bounded by $\tilde{y}_1$ and $\tilde{y}_2$.*

Proof. Denote the oriented paths connecting $\tilde{y}_1$ to $\tilde{y}_2$ and $\tilde{f}(\tilde{y}_1)$ to $\tilde{f}(\tilde{y}_2)$ by $\tilde{\alpha}_0$ and $\tilde{\alpha}_1$ respectively. Let $r : \Gamma \rightarrow \tilde{\alpha}_1$ be retraction onto the nearest point in $\tilde{\alpha}_1$ and let $g = r\tilde{f} : \tilde{\alpha}_0 \rightarrow \tilde{\alpha}_1$. By hypothesis, $\tilde{\alpha}_0$ is a proper subpath of $\tilde{\alpha}_1$ and g is a surjection. If $\tilde{y}$ is the first point in $\tilde{\alpha}_0$ such that $\tilde{g}(\tilde{y}) = \tilde{y}$ then $\tilde{y}_1 < \tilde{y} < \tilde{y}_2$ and $\tilde{g}(\tilde{z}) < \tilde{g}(\tilde{y})$ for $y_1 < z < y$. It follows that $\tilde{f}(\tilde{y}) \in \tilde{\alpha}_1$ and hence that $\tilde{y}$ is fixed by $\tilde{f}$. $\square$

Corollary 3.17. *If $\tilde{f}$ is a principal lift then $\text{Fix}(\tilde{f}) \neq \emptyset$.*

Proof. Suppose that there is a non-trivial covering translation T_c that has its endpoints in $\text{Fix}_N(\hat{f})$ and so commutes with $\tilde{f}$. Assuming without loss that A_c is fixed point free, there is a point in A_c that moves toward one of the endpoints of A_c , say P . Since $\tilde{f}$ commutes with T_c , there are points in Γ that are arbitrarily close to P and that move toward P . The same property holds for an attractor $P \in \text{Fix}(\hat{f})$ by Lemma 3.15. One may therefore choose distinct $P_1, P_2 \in \text{Fix}_N(\hat{f})$ and $\tilde{x}_1, \tilde{x}_2 \in \tilde{\Gamma}$ such that $\tilde{x}_i$ is close to and moves toward P_i . It follows that $\tilde{x}_1$ and $\tilde{x}_2$ move away from each other. Lemma 3.16 produces the desired fixed point. $\square$

There are two cases in which a lift $\tilde{f}$ corresponding to a Nielsen class in $\text{Fix}(f)$ is not principal. The first arises from a ‘non-singular’ leaf of an attracting lamination as noted in Remark 3.6; in this case $\text{Fix}(\tilde{f})$ is a single point. In the second case, there is a circle component of $\text{Fix}(f)$ with no outward pointing periodic directions and $\text{Fix}(\tilde{f})$ is an axis A_c . The second type could be eliminated by adding properties to Theorem 2.19. We allow the circle components for now and defer the additional properties until section 4.

Periodic points for f are *Nielsen equivalent* if they are Nielsen equivalent as fixed points for some iterate of f .

Definition 3.18. We say that x is *principal* if neither of the following conditions are satisfied.

- x is the only element of $\text{Per}(f)$ in its Nielsen class and there are exactly two periodic directions at x , both of which are contained in the same EG stratum.
- x is contained in a component C of $\text{Per}(f)$ that is topologically a circle and each point in C has exactly two periodic directions.

Lifts to Γ of principal periodic points in G are said to be *principal*. If each principal vertex and each periodic direction at a principal vertex has period one then we say that $f : G \rightarrow G$ is *rotationless*.

In practice, we only apply these definitions to $f : G \rightarrow G$ that satisfy the conclusions of Theorem 2.19. In particular, by item (4) of Lemma 2.20, there are at least two periodic directions at each $x \in \text{Per}(f)$.

Principal periodic points are either contained in periodic edges or are vertices. Thus every $f : G \rightarrow G$ has a rotationless iterate. Any endpoint of an indivisible periodic Nielsen path is principal as is the initial endpoint of any non-periodic NEG edge. The latter implies that each NEG stratum in a rotationless relative train track map is a single edge.

The following lemma shows that an EG stratum has at least one principal vertex.

Lemma 3.19. *Assume that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19. For every EG stratum H_r there is a principal vertex whose link contains a periodic direction in H_r .*

Proof. If some vertex $v \in H_r$ belongs to a non-contractible component of G_{r-1} then v is periodic and there is at least one periodic direction in H_r by Lemma 2.9. There is also at least one periodic direction at v determined by an edge of G_{r-1} so v is principal. If there is no such vertex, then H_r is a union of components of G_r . Lemma 5.2 of [3] states there is a principal lift of some iterate of $f|H_r$. In the course of proving this lemma, it is shown that either there is a vertex with three periodic directions or there is an indivisible periodic Nielsen path in H_r . In either case there is a vertex that is principal for $f|H_r$ and hence also for f . $\square$

Remark 3.20. Lemma 3.19 implies that the transition matrix M_r of an EG stratum of a rotationless $f : G \rightarrow G$ satisfying the conclusions of Theorem 2.19 has at least one non-zero diagonal entry and so is aperiodic. For each $\Lambda \in \mathcal{L}(\phi)$ there is an EG stratum H_r such that Λ has height r and this defines a bijection (see Definition 3.1.12 of [2]) between $\mathcal{L}(\phi)$ and the set of EG strata.

The next lemma relates an attractor in $\text{Fix}_N(\hat{f})$ to a fixed direction of Df .

Lemma 3.21. *Suppose that $\tilde{f}$ is a principal lift of a relative train track map $f : G \rightarrow G$.*

- (1) *For each attractor $P \in \text{Fix}_N(\hat{f})$ there is a (not necessarily unique) $\tilde{x} \in \text{Fix}(\tilde{f})$ such that the interior of the ray $\tilde{R}_{\tilde{x},P}$ that starts at $\tilde{x}$ and that converges to P is fixed point free.*
- (2) *If $P \in \text{Fix}_N(\hat{f})$ is an attractor, if $\tilde{x} \in \text{Fix}(\tilde{f})$ and if the interior of $\tilde{R}_{\tilde{x},P}$ is fixed point free then no point in the interior of $\tilde{R}_{\tilde{x},P}$ is mapped by $\tilde{f}$ to $\tilde{x}$; in particular, the initial direction determined by $\tilde{R}_{\tilde{x},P}$ is fixed.*
- (3) *If P and Q are distinct attractors in $\text{Fix}_N(\hat{f})$, if $\tilde{x} \in \text{Fix}(\tilde{f})$ and if the interiors of both $\tilde{R}_{\tilde{x},P}$ and $\tilde{R}_{\tilde{x},Q}$ are fixed point free then the directions determined by $\tilde{R}_{\tilde{x},P}$ and $\tilde{R}_{\tilde{x},Q}$ are distinct.*

Proof. To find $\tilde{x} \in \text{Fix}(\tilde{f})$ and $\tilde{R}_{\tilde{x},P}$ as in (1), start with any ray $\tilde{R}'$ whose initial point is in $\text{Fix}(\tilde{f})$ and that converges to P and let $\tilde{R}_{\tilde{x},P}$ be the subray of $\tilde{R}'$ that begins at the last point $\tilde{x}$ of $\text{Fix}(\tilde{f})$ in $\tilde{R}'$. If $\tilde{R}_{\tilde{x},P}$ and $\tilde{R}_{\tilde{x},Q}$ are as in (3) and have the same initial edge then their ‘difference’ would be a fixed point free line whose ends converge to attractors in contradiction to Lemma 3.15 and Lemma 3.16. This verifies (3). By the same reasoning, no points in the interior of $\tilde{R}_{\tilde{x},P}$ can map to $\tilde{x}$, which implies that the initial edge of $\tilde{R}_{\tilde{x},P}$ determines a fixed direction at $\tilde{x}$. This proves (2). $\square$

Corollary 3.22. *Assume that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19. If $\tilde{f}$ is a principal lift then each element of $\text{Fix}(\tilde{f})$ is principal.*

Proof. Let Φ be the automorphism corresponding to $\tilde{f}$. If $\text{Fix}(\Phi)$ has rank at least two then $\text{Fix}(\tilde{f})$ is neither a single point nor a single axis and we are done (see Corollary 3.17 and Lemma 2.1). If $\text{Fix}(\Phi)$ has rank one then $\text{Fix}(\tilde{f})$ is infinite and

$\text{Fix}_N(\hat{f})$ contains an attractor by the definition of principal lift and by Lemma 2.3. Lemma 3.21 implies that some $\tilde{x} \in \text{Fix}(\tilde{f})$ has a fixed direction that does not come from a fixed edge and again we are done. In the remaining case, $\text{Fix}(\hat{f})$ is a finite set of attractors and does not contain the endpoints of any axis. Obviously $\text{Fix}(\tilde{f})$ is not an axis. Suppose that $\text{Fix}(\tilde{f})$ is a single point $\tilde{x}$, that there are only two periodic directions at $\tilde{x}$ and these two directions are determined by lifts $\tilde{E}_1$ and $\tilde{E}_2$ of oriented edges of the same EG stratum H_r . Lemma 3.21 and Lemma 2.13 imply that $\text{Fix}(\hat{f})$ is the endpoint set of a generic leaf of an element of $\mathcal{L}(\phi)$ in contradiction to the assumption that $\tilde{f}$ is principal. We conclude that $\tilde{x}$ is principal as desired. $\square$

To prove the converse we use fixed directions of $\tilde{f}$ to find elements of $\text{Fix}_N(\hat{f})$. The following lemma is from [2]; the proof is short and is repeated for the readers convenience.

Lemma 3.23. *If $\text{Fix}(\tilde{f}) = \emptyset$ then there is a ray $\tilde{R} \subset \Gamma$ converging to an element $P \in \text{Fix}(\hat{f})$ and there are points in $\tilde{R}$ arbitrarily close to P that move toward P .*

Proof. For each vertex $\tilde{y}$ of Γ , we say that the initial edge of the path from $\tilde{y}$ to $h(\tilde{y})$ is *preferred* by $\tilde{y}$. Starting with any vertex $\tilde{y}_0$, inductively define $\tilde{y}_{i+1}$ to be the other endpoint of the edge preferred by $\tilde{y}_i$. If $\tilde{E}$ is preferred by both of its endpoints then $\tilde{f}$ maps a proper subinterval of $\tilde{E}$ over all of $\tilde{E}$ (reversing orientation) in contradiction to the assumption that $\text{Fix}(\tilde{f}) = \emptyset$. It follows that the $\tilde{y}_i$'s are contained in a ray that converges to some $P \in \text{Fix}(\hat{f})$ and that $\tilde{y}_i$ moves toward P . $\square$

We isolate the following notation and lemma for reference throughout the paper.

Notation 3.24. Suppose that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19, that H_r is a single edge E_r and that $f(E_r) = E_r u$ for some non-trivial path $u \subset G_{r-1}$. Let $\tilde{E}_r$ be a lift of E_r and let $\tilde{f} : \Gamma \rightarrow \Gamma$ be the lift of f that fixes the initial endpoint of $\tilde{E}_r$. By (NEG), the component C of G_{r-1} that contains the terminal endpoint w of E_r is not contractible. Denote the copy of the universal cover of C that contains the terminal endpoint of $\tilde{E}_r$ by Γ_{r-1} and the restriction $\tilde{f}|_{\Gamma_{r-1}}$ by $h : \Gamma_{r-1} \rightarrow \Gamma_{r-1}$.

The covering translations that preserve Γ_{r-1} define a free factor $F(C)$ of F_n such that $[[F(C)]] = [[\pi_1(C)]]$. The closure in ∂F_n of $\{T_c^\pm : c \in F(C)\}$ is naturally identified with $\partial F(C)$ and with the space of ends of Γ_{r-1} . Moreover, $\hat{h} = \hat{f}|_{\partial F(C)} : \partial F(C) \rightarrow \partial F(C)$.

Lemma 3.25. *Assume that $\tilde{f}$ and h are as in Notation 3.24. If $\text{Fix}(h) = \emptyset$ then there is a ray $\tilde{R} \subset \Gamma_{r-1}$ converging to an element $P \in \text{Fix}(\hat{h})$ and there are points in $\tilde{R}$ arbitrarily close to P that move toward P .*

Proof. This follows from Lemma 3.23 applied to $h : \Gamma_{r-1} \rightarrow \Gamma_{r-1}$. $\square$

Our next result is an extension of Lemma 2.13.

Lemma 3.26. *Suppose that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19 and is rotationless, that $\tilde{f} : \Gamma \rightarrow \Gamma$ is a lift of f , that $\tilde{v} \in \text{Fix}(\tilde{f})$ and that $D\tilde{f}$ fixes the direction at $\tilde{v}$ determined by a lift $\tilde{E}$ of an edge $E \subset H_r$. Then there exists $P \in \text{Fix}(\hat{f})$ so that the ray $\tilde{R}$ from the initial endpoint of $\tilde{E}$ to P contains $\tilde{E}$ and satisfies the following properties.*

- (1) *There are points in $\tilde{R}$ arbitrarily close to P that are either fixed or move toward P . If there does not exist $c \in F_n$ such that $\text{Fix}(\hat{f}) = \{T_c^\pm\}$ then $P \in \text{Fix}_N(\hat{f})$.*
- (2) *If H_r is EG then P is an attractor whose accumulation set is the unique attracting lamination of height r , the interior of $\tilde{R}$ is fixed point free and $\tilde{R}$ projects to an r -legal ray in G_r .*
- (3) *If H_r is NEG and non-fixed then $\tilde{R} \setminus \tilde{E}$ projects into G_{r-1} .*
- (4) *No point in the interior of $\tilde{R}$ is mapped to $\tilde{v}$ by any iterate of $\tilde{f}$.*

Proof. The second part of (1) follows from the first part of (1) and Lemma 3.15.

The proof is by induction on r , starting with $r = 1$. If $G_1 \subset \text{Fix}(f)$ then we may choose P to be the endpoint of any ray $\tilde{R}$ that begins with $\tilde{E}$ and projects into G_1 ; the existence of such a ray follows from the fact (Theorem 2.19(F)) that G_1 is its own core. If G_1 is EG then the existence of P follows from Lemma 2.13 and Lemma 3.21. This completes the $r = 1$ case so we may now assume that the lemma holds for edges with height less than r .

If H_r is EG then the existence of P follows from Lemma 2.13 and Lemma 3.21. We may therefore assume that H_r is NEG. Let $h : \Gamma_{r-1} \rightarrow \Gamma_{r-1}$ be as in Notation 3.24. If $\text{Fix}(h) \neq \emptyset$, then the initial endpoint of $\tilde{E}$ and some $\tilde{x} \in \text{Fix}(h)$ cobound an indivisible Nielsen path. Thus $\tilde{x}$ is principal, there is a fixed direction in Γ_{r-1} at $\tilde{x}$ and the existence of an appropriate $P \in \text{Fix}(\hat{h})$ follows from the inductive hypothesis. The case that $\text{Fix}(h) = \emptyset$ follows from Lemma 3.25. $\square$

We now can prove the converse to Corollary 3.22 under the assumption that $f : G \rightarrow G$ is rotationless.

Corollary 3.27. *Suppose that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19 and is rotationless. If some, and hence every, $\tilde{x} \in \text{Fix}(\tilde{f})$ is principal then $\tilde{f}$ is principal.*

Proof. Assume that $\text{Fix}(\tilde{f})$ consists of principal points. Since $f : G \rightarrow G$ is rotationless, periodic directions based in $\text{Fix}(\tilde{f})$ are fixed. Lemma 2.20(4) implies that each $\tilde{x} \in \text{Fix}(\tilde{f})$ has at least two fixed directions. If some $\tilde{x} \in \text{Fix}(\tilde{f})$ has at least three fixed directions, then Lemma 3.26 produces at least three points in $\text{Fix}_N(\hat{f})$ and we are done. We may therefore assume that there are exactly two fixed, and hence exactly two periodic, directions at each $\tilde{x} \in \text{Fix}(\tilde{f})$. If $\text{Fix}(\tilde{f})$ contains an edge, then by Definition 3.18 there must be such an edge with a valence one vertex in $\text{Fix}(f)$.

This contradicts items (1) and (4) of Lemma 2.20 and we conclude that there are no fixed edges. Choose an edge $E \subset H_r$ and a lift $\tilde{E}$ whose initial direction is fixed and based at some $\tilde{x} \in \text{Fix}(\tilde{f})$. Let $\tilde{R}$ be the ray that begins with $\tilde{E}$ and ends at some $P \in \text{Fix}(\hat{f})$ as in Lemma 3.26.

If H_r is EG then the accumulation set of P is an attracting lamination which implies by Lemma 3.1.16 of [2] that P is not the endpoint of an axis. If H_r is NEG then the accumulation set of P is contained in G_{r-1} which implies that P is not the endpoint of an axis that contains $\tilde{E}$. It follows that the line composed of $\tilde{R}$ and the ray determined by the second fixed direction at $\tilde{x}$ is not an axis. We have now shown that $\text{Fix}(\hat{f})$ is not the endpoint set of an axis and hence that every point in $\text{Fix}(\hat{f})$ produced by Lemma 3.26 is contained in $\text{Fix}_N(\hat{f})$. Thus $\text{Fix}_N(\hat{f})$ contains at least two points and is not the endpoint set of an axis.

To complete the proof we assume that $\text{Fix}_N(\hat{f})$ is the endpoint set of a lift $\tilde{\ell}$ of a generic leaf of an attracting lamination and argue to a contradiction. Since ℓ is birecurrent and contains E , H_r is EG and the second fixed direction based at $\tilde{x}$ comes from an edge in H_r . Lemma 3.26(2) implies that $\ell \subset G_r$ is r -legal and hence does not contain any indivisible Nielsen paths of height r . But then $\tilde{x}$ must be the only fixed point in $\tilde{\ell}$. Since $\text{Fix}(\tilde{f})$ is principal it must contain a point other than $\tilde{x}$ and that point would have a fixed direction that does not come from the initial edge of a ray converging to an endpoint of $\tilde{\ell}$. This contradiction completes the proof. $\square$

3.4 Rotationless is Rotationless

We prove in this section that rotationless relative train track maps represent forward rotationless outer automorphisms and vice-versa.

Lemma 3.28. *Suppose that $f : G \rightarrow G$ satisfies the conclusions of Theorem 2.19 and is rotationless. Every periodic Nielsen path σ with principal endpoints has period one.*

Proof. There is no loss in assuming that σ is either a single edge or an indivisible periodic Nielsen path. In the former case, σ is a periodic edge with an principal endpoint and so is fixed. We may therefore assume that σ is indivisible.

The proof is by induction on the height r of σ with the $r = 0$ case being vacuously true. Let p be the period of σ and let $v \in \text{Fix}(f)$ be an endpoint of σ . The case that H_r is EG follows from Lemma 2.11(3).

We may therefore assume that H_r is a single non-fixed NEG edge E_r . Lemma 4.1.4 of [2] implies that after reversing the orientation on σ if necessary, $\sigma = E_r\mu$ or $\sigma = E_r\mu\bar{E}_r$ for some path $\mu \subset G_{r-1}$. Let $\tilde{E}_r$ be a lift of E_r with initial endpoint $\tilde{v}$, let $\tilde{f}$ be the lift that fixes $\tilde{v}$ and let $h : \Gamma_{r-1} \rightarrow \Gamma_{r-1}$ be as in Notation 3.24. By Lemma 3.27, $\tilde{f}$ is principal. Denote the terminal endpoint of the lift $\tilde{\sigma}$ that begins at $\tilde{v}$ by $\tilde{w}$.

If $\sigma = E_r\mu$ then $\tilde{w} \in \Gamma_{r-1}$. If $p \neq 1$ then the path $\tilde{\tau}$ connecting $\tilde{w}$ to $h(\tilde{w})$ projects to a non-trivial periodic Nielsen path $\tau \subset G_{r-1}$ that is closed because $\tilde{w}$ projects to $w \in \text{Fix}(f)$. Since $\tilde{w}$ is principal, the inductive hypothesis implies that τ has period one and hence that the projection of the closed path $\tilde{\tau}h(\tilde{\tau}) \dots h^{p-1}(\tilde{\tau})$ to G_{r-1} is homotopic to τ^p . This contradicts the fact that τ and hence τ^p determines a non-trivial conjugacy class in F_n . Thus $p = 1$ in the case that $\sigma = E_r\mu$.

Suppose now that $\sigma = E_r\mu\tilde{E}_r$. If $\text{Fix}(h^p) \neq \emptyset$ then the path $\tilde{\sigma}_1$ connecting $\tilde{v}$ to $\tilde{x} \in \text{Fix}(h^p)$ and the path $\tilde{\sigma}_2$ connecting $\tilde{x}$ to $\tilde{w}$ are periodic Nielsen paths. By the preceding case σ_1 and σ_2 , and hence σ , has period one. We may therefore assume that $\text{Fix}(h^p) = \emptyset$.

Let $T_c : \Gamma \rightarrow \Gamma$ be the covering translation satisfying $T_c(\tilde{v}) = \tilde{w}$. Then T_c commutes with $\tilde{f}^p$, the axis A_c is contained in Γ_{r-1} and $T_c^\pm \in \text{Fix}(\hat{h}^p)$. If Φ is the principal automorphism corresponding to $\tilde{f}$ then $\hat{T}_{\Phi(c)} = \tilde{f}T_c\tilde{f}^{-1}$, which implies that $T_{\Phi(c)}^\pm = \hat{h}(T_c^\pm) \in \text{Fix}(\hat{h}^p)$ and that $A_{\Phi(c)} \subset \Gamma_{r-1}$. If $\{T_c^\pm\}$ is not $\hat{h}$ -invariant, then $\text{Fix}_N(\hat{h}^p)$ contains the four points $\{T_c^\pm\} \cup \{\hat{h}(T_c^\pm)\}$ and h^p is a principal lift of $f|_C$ where C is the component of G_{r-1} that contains the terminal endpoint of E_r . This contradicts Corollary 3.17 and the assumption that $\text{Fix}(h^p) = \emptyset$. Thus $\{T_c^\pm\}$ is $\hat{h}$ -invariant. If $\hat{h}$ interchanges $T_c^\pm$ then $\text{Fix}_N(\hat{h}^p)$ contains $\{T_c^\pm\}$ and at least one point in $\text{Fix}(\hat{h})$ by Lemma 3.25. This contradicts Corollary 3.17 and we conclude that $T_c^\pm \in \text{Fix}(\hat{h})$. It follows that $\tilde{f}$ commutes with T_c and hence that $\tilde{w} \in \text{Fix}(\tilde{f})$. This proves that $p = 1$ and so completes the inductive step. $\square$

Proposition 3.29. *Suppose that $f : G \rightarrow G$ represents ϕ and satisfies the conclusions of Theorem 2.19. Then $f : G \rightarrow G$ is rotationless if and only if ϕ is forward rotationless.*

Proof. Suppose that $f : G \rightarrow G$ is rotationless, that $k \geq 1$ and that $\tilde{g} : \Gamma \rightarrow \Gamma$ is a principal lift of $g := f^k$. Corollary 3.17 and Corollary 3.22 imply that $\text{Fix}(\tilde{g})$ is a non-empty set of principal fixed points. Since f is rotationless, for each $\tilde{v} \in \text{Fix}(\tilde{g})$ there is a lift $\tilde{f} : \Gamma \rightarrow \Gamma$ that fixes $\tilde{v}$ and all periodic directions at $\tilde{v}$. To prove that ϕ is forward rotationless it suffices to show that $\text{Fix}_N(\hat{f}) = \text{Fix}_N(\hat{g})$ and hence (Remark 3.4) that $\tilde{f}^k = \tilde{g}$.

The path connecting $\tilde{v}$ to another point in $\text{Fix}(\tilde{g})$ projects to a Nielsen path for g and hence by Lemma 3.28, a Nielsen path for f . Thus $\text{Fix}(\tilde{f}) = \text{Fix}(\tilde{g})$. It follows that $\tilde{g}$ and $\tilde{f}$ commute with the same covering translations and Lemma 2.3 implies that $\text{Fix}_N(\hat{f})$ and $\text{Fix}_N(\hat{g})$ have the same non-isolated points.

Each isolated point $P \in \text{Fix}_N(\hat{g})$ is an attractor for $\hat{g}$. It suffices to show that $P \in \text{Fix}_N(\hat{f})$. By Lemma 3.21 there is a ray $\tilde{R}$ that terminates at P , that intersects $\text{Fix}(\tilde{g})$ only in its initial endpoint and whose initial direction is fixed by $D\tilde{g}$, and hence also by $D\tilde{f}$. We may assume that the height r of the initial edge $\tilde{E}$ of $\tilde{R}$ is minimal among all choices of $\tilde{R}$. By Lemma 3.26, $\tilde{E}$ extends to a ray $\tilde{R}'$ that converges to some $P' \in \text{Fix}(\hat{f})$. It suffices to show that $P' = P$ since a repeller for $\hat{f}$ could not be an attractor for $\hat{g}$. If H_r is EG this follows from Lemma 3.26(2) and Lemma 3.21(3) applied to g . We may therefore assume that H_r is NEG. If there exists $\tilde{x} \in \text{Fix}(\tilde{g}) \cap \tilde{R}'$

then $\tilde{x} \in \tilde{R}' \setminus \tilde{E}$ and the ray connecting $\tilde{x}$ to P is contained in G_{r-1} in contradiction to our choice of r . We may therefore assume that $\text{Fix}(\tilde{g}) \cap \tilde{R}' = \emptyset$. Lemma 3.26(1) implies that there exists $\tilde{x} \in \tilde{R}'$ that is moved toward P' by $\tilde{f}$ and Lemma 3.16 then implies that $P = P'$. This completes the proof of the only if direction of the proposition.

For the if direction, assume that ϕ is forward rotationless and choose $k > 0$ so that $g := f^k$ is rotationless. For each principal $v \in \text{Fix}(g)$, there exist a lift $\tilde{v}$ of v and a principal lift $\tilde{g}$ of g that fixes $\tilde{v}$. Since ϕ is forward rotationless, there is a lift $\tilde{f}$ of f such that $\tilde{f}^k = \tilde{g}$ and such that $\text{Fix}_N(\tilde{f}) = \text{Fix}_N(\tilde{g})$. It suffices to show that $\tilde{v} \in \text{Fix}(\tilde{f})$ and that each $D\tilde{g}$ -fixed direction $\tilde{d}_1$ at $\tilde{v}$ is $D\tilde{f}$ -fixed.

The edge determined by $\tilde{d}_1$ extends to a ray $\tilde{R}_1$ that converges to some $P_1 \in \text{Fix}_N(\tilde{g}) = \text{Fix}_N(\tilde{f})$. Define P_2 and $\tilde{R}_2$ similarly using a second $D\tilde{g}$ -fixed direction $\tilde{d}_2$ based at $\tilde{v}$ and denote the line connecting P_1 to P_2 by $\tilde{\gamma}$. Thus $\tilde{f}_\#(\tilde{\gamma}) = \tilde{\gamma}$ and the turn $(\tilde{d}_1, \tilde{d}_2)$ is legal for $\tilde{g}$ and hence for $\tilde{f}$. If $\tilde{f}(\tilde{v}) \notin \tilde{\gamma}$ then there exists $\tilde{y} \in \tilde{\gamma}$ not equal to $\tilde{v}$ such that $\tilde{f}(\tilde{y}) = \tilde{f}(\tilde{v})$. But then $\tilde{f}^k(\tilde{y}) = \tilde{f}^k(\tilde{v}) = \tilde{v}$ which contradicts Lemma 3.26(4) applied to $\tilde{g}$. This proves that $\tilde{f}(\tilde{v}) \in \tilde{\gamma}$. Suppose that $\tilde{f}(\tilde{v}) \neq \tilde{v}$. Denote $\tilde{v}$ by $\tilde{v}_0$ and orient $\tilde{\gamma}$ so that P_1 is the negative end and P_2 is the positive end. Assuming without loss that $\tilde{v} < \tilde{f}(\tilde{v})$ in the order induced from the orientation, there exist $\tilde{v}_i \in \tilde{\gamma}$ for $1 \leq i \leq k$ such that $\tilde{v}_i < \tilde{v}_{i-1}$ and such that $\tilde{f}(\tilde{v}_i) = \tilde{v}_{i-1}$. But then $\tilde{f}^k(\tilde{v}_k) = \tilde{v}$ in contradiction to Lemma 3.26(4). We conclude that $\tilde{f}(\tilde{v}) = \tilde{v}$. A third application of Lemma 3.26(4) implies that the directions $\tilde{d}_i$ are fixed by $D\tilde{f}$. $\square$

3.5 Properties of Forward Rotationless ϕ

Lemma 3.30. *The following hold for each forward rotationless $\phi \in \text{Out}(F_n)$.*

- (1) *Each periodic conjugacy class is fixed and each representative of that conjugacy class is fixed by some principal automorphism representing ϕ .*
- (2) *Each $\Lambda \in \mathcal{L}(\phi)$ is ϕ -invariant.*
- (3) *A free factor that is invariant under an iterate of ϕ is ϕ -invariant.*

Proof. If the conjugacy class of c is fixed by ϕ^k for $k \geq 1$ then by Remark 3.2 there exists a principal automorphism $\Phi_k \in P(\phi^k)$ that fixes c . By Lemma 2.1 this is equivalent to $T_c^\pm \in \text{Fix}_N(\hat{\Phi}_k)$. Since ϕ is forward rotationless, we may assume that $k = 1$. This completes the proof of the first item.

If $f : G \rightarrow G$ is a rotationless relative train track map that represents ϕ and that satisfies the conclusions of Theorem 2.19 then each $\Lambda \in \mathcal{L}(\phi)$ is associated to a unique EG stratum H_r ; see Definition 3.1.12 of [2] for details. Lemma 3.19 and Lemma 3.26(2) imply that Λ is the accumulation set of an attractor $P \in \text{Fix}_N(\hat{\Phi})$ for some principal automorphism representing ϕ . This implies (2).

For the third item, suppose that the free factor F is ϕ^k -invariant for some $k \geq 1$. If F has rank one then it is ϕ -invariant by the first item of this lemma. We may

therefore assume that F has rank at least two. Let $\mathcal{C}$ be the set of bi-infinite lines γ that are carried by F and for which there exist a principal lift Φ of an iterate of ϕ and a lift $\tilde{\gamma}$ of γ whose endpoints are contained in $\text{Fix}_N(\hat{\Phi})$. Since ϕ is forward rotationless, each γ is ϕ -invariant, so $\mathcal{C}$ is ϕ -invariant. Obviously $\mathcal{C}$ is carried by F so to prove that F is ϕ -invariant it suffices, by Corollary 2.5, to show that no proper ϕ -invariant free factor system $\mathcal{F}$ of F carries $\mathcal{C}$.

Suppose to the contrary that such an $\mathcal{F}$ exists. By Theorem 2.19 there is a relative train track map $g : G' \rightarrow G'$ representing $\phi^k|F$ in which $\mathcal{F}$ is represented by a proper filtration element $G'_r \subset G'$. After replacing $\phi^k|F$ and f by iterates we may assume that they are (forward) rotationless. There is a principal vertex $v \in G'$ whose link contains an edge E of $G' \setminus G'_r$ that determines a fixed direction. This follows from Lemma 3.19 if there is an EG stratum in $G' \setminus G'_r$ and from the definition of principal otherwise. Lemma 3.26 and the fact that there are at least two periodic directions based at v imply that there is a principal lift $\tilde{f}' : \Gamma' \rightarrow \Gamma'$ and a line $\tilde{\gamma}$ whose endpoints are contained in $\text{Fix}_N(\hat{f}')$ and whose projected image γ crosses E and so is not carried by G'_r . The automorphism $\Phi' \in P(\phi|F)$ determined by $\tilde{f}'$ extends to an element $\Phi \in P(\phi^k)$ with $\text{Fix}_N(\hat{\Phi}') \subset \text{Fix}_N(\hat{\Phi})$. Thus $\gamma \in \mathcal{C}$ in contradiction to our choice of $\mathcal{F}$ and G'_r . $\square$

Corollary 3.31. *If ϕ is forward rotationless and F is a ϕ -invariant free factor, then $\theta := \phi|F \in \text{Out}(F)$ is forward rotationless.*

Proof. Lemma 3.30(1) handles the case that F has rank one so we may assume that F has rank at least two. Choose a relative train track map $f : G \rightarrow G$ and filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ satisfying the conclusions of Theorem 2.19 and representing ϕ such that the conjugacy class of F is represented by G_l for some l . Proposition 3.29 implies that $f : G \rightarrow G$ is rotationless. The restriction of $f : G \rightarrow G$ to G_l is a rotationless relative train track map representing θ and satisfying the conclusions of Theorem 2.19. A second application of Proposition 3.29 implies that θ is forward rotationless. $\square$

4 Completely Split Relative Train Track Maps

For every $\phi \in \text{Out}(F_n)$ there exists $k > 0$ such that ϕ^k is represented by an improved relative train track map (IRT) $f : G \rightarrow G$ as defined in Theorem 5.1.5 of [2]. In this section we update this theorem, replacing IRTs with CTs, by controlling the iteration index k , adding a very useful property called complete splitting, and by making small changes to previous definitions. Section 4.1 contains all the necessary definitions. In section 4.2 we show that completely splittings in a CT are hard splittings in the sense of [5]. A detailed comparison of IRTs and CTs is given in section 4.3. There is one new move needed for the construction of CTs. It is defined in section 4.4 and the existence theorem is stated and proved in section 4.5. A few additional properties of CTs are presented in section 4.6

4.1 Definitions and Notation

For $a \in F_n$, we let $[a]_u$ be the *unoriented conjugacy class determined by a* . Thus, $[a]_u = [b]_u$ if and only if a is conjugate to either of a or $\bar{a}$. If σ is a closed path then we let $[\sigma]_u$ be the *unoriented conjugacy class determined by σ* , thought of as a circuit.

Suppose that $f : G \rightarrow G$ is a rotationless relative train track map with filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$. Each NEG stratum H_i is a single edge E_i satisfying $f(E_i) = E_i u_i$ for some (necessarily closed by (NEG)) path $u_i \subset G_{i-1}$ that is sometimes called the *suffix* for E_i . If u_i is a non-trivial Nielsen path, then we say that E_i is a *linear edge*. In the linear case, we define the *axis* or *twistor* for E_i to be $[w_i]_u$ where w_i is root-free and $u_i = w_i^{d_i}$ for some $d_i \neq 0$.

Definition 4.1. If E_i and E_j are linear edges and if there are $m_i, m_j > 0$ and a closed root-free Nielsen path w such that $u_i = w^{m_i}$ and $u_j = w^{m_j}$ then a path of the form $E_i w^p \bar{E}_j$ with $p \in \mathbb{Z}$ is called an *exceptional path*.

Remark 4.2. If $E_i w^p \bar{E}_j$ is an exceptional path then $f_{\#}^k(E_i w^p \bar{E}_j) = E_i w^{p+k(m_i-m_j)} \bar{E}_j$ for all $k \geq 0$. It follows that $E_i w^p \bar{E}_j$ is a Nielsen path if and only if $m_i = m_j$, that $f_{\#}$ induces a height preserving bijection on the set of exceptional paths and that the interior of $E_i w^p \bar{E}_j$ is an increasing union of pre-trivial paths.

Definition 4.3. A filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ that satisfies the following property is said to be *reduced (with respect to ϕ)*: if a free factor system $\mathcal{F}'$ is ϕ^k -invariant for some $k > 0$ and if $\mathcal{F}(G_{r-1}) \sqsubset \mathcal{F}' \sqsubset \mathcal{F}(G_r)$ then either $\mathcal{F}' = \mathcal{F}(G_{r-1})$ or $\mathcal{F}' = \mathcal{F}(G_r)$.

Definition 4.4. If E is an edge in an irreducible stratum H_r and $k > 0$ then a maximal subpath σ of $f_{\#}^k(E)$ in a zero stratum H_i is said to be *r -taken* or just *taken* if r is irrelevant. Note that if H_i is enveloped by an EG stratum H_s then σ has endpoints in H_s and so is a connecting path. A non-trivial path or circuit σ is *completely split* if it has a splitting, called a *complete splitting*, into subpaths, each of which is either a single edge in an irreducible stratum, an indivisible Nielsen path, an exceptional path or a taken connecting path in a zero stratum.

Definition 4.5. A relative train track map is *completely split* if

- (1) $f(E)$ is completely split for each edge E in each irreducible stratum.
- (2) If σ is a taken connecting path in a zero stratum then $f_{\#}(\sigma)$ is completely split.

The next lemma states that if $f : G \rightarrow G$ is completely split then $f_{\#}$ maps completely split paths to completely split paths.

Lemma 4.6. *If $f : G \rightarrow G$ is completely split and σ is a completely split path or circuit then $f_{\#}(\sigma)$ is completely split. Moreover if $\sigma = \sigma_1 \cdot \dots \cdot \sigma_k$ is a complete splitting then $f_{\#}(\sigma)$ has a complete splitting that refines $f_{\#}(\sigma) = f_{\#}(\sigma_1) \cdot \dots \cdot f_{\#}(\sigma_s)$*

Proof. This is immediate from the definitions and the fact that $f_{\#}$ carries indivisible Nielsen paths to indivisible Nielsen paths and exceptional paths to exceptional paths. $\square$

We now come to our main definition. When equivalent descriptions of a property are available, for example in (EG Nielsen Paths), we have chosen the one that is easiest to check.

Definition 4.7. A relative train track map $f : G \rightarrow G$ and filtration $\mathcal{F}$ given by $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$ is said to be a *CT* (for completely split improved relative train track map) if it satisfies the following properties.

- (1) **(Rotationless)** $f : G \rightarrow G$ is rotationless. (See Remark 4.8.)
- (2) **(Completely Split)** $f : G \rightarrow G$ is completely split.
- (3) **(Filtration)** $\mathcal{F}$ is reduced. The core of each filtration element is a filtration element.
- (4) **(Vertices)** The endpoints of all indivisible periodic (necessarily fixed) Nielsen paths are (necessarily principal) vertices. The terminal endpoint of each non-fixed NEG edge is principal (and hence fixed). (See Remark 4.9 and Lemma 4.22.)
- (5) **(Periodic Edges)** Each periodic edge is fixed and each endpoint of a fixed edge is principal. If the unique edge E_r in a fixed stratum H_r is not a loop then G_{r-1} is a core graph and both ends of E_r are contained in G_{r-1} .
- (6) **(Zero Strata)** If H_i is a zero stratum, then H_i is enveloped by an EG stratum H_r , each edge in H_i is r -taken and each vertex in H_i is contained in H_r and has link contained in $H_i \cup H_r$.
- (7) **(Linear Edges)** For each linear E_i there is a closed root-free Nielsen path w_i such that $f(E_i) = E_i w_i^{d_i}$ for some $d_i \neq 0$. If E_i and E_j are distinct linear edges with the same axes then $w_i = w_j$ and $d_i \neq d_j$. (See Remark 4.11.)
- (8) **(NEG Nielsen Paths)** If the highest edges in an indivisible Nielsen path σ belong to an NEG stratum then there is a linear edge E_i with w_i as in (Linear Edges) and there exists $k \neq 0$ such that $\sigma = E_i w_i^k \bar{E}_i$.
- (9) **(EG Nielsen Paths)** (See also Lemmas 4.18 and 4.19 and Corollaries 4.20 and 4.34) If H_r is EG and ρ is an indivisible Nielsen path of height r , then $f|_{G_r} = \theta \circ f_{r-1} \circ f_r$ where :
 - (a) $f_r : G_r \rightarrow G^1$ is a composition of proper extended folds defined by iteratively folding ρ .
 - (b) $f_{r-1} : G^1 \rightarrow G^2$ is a composition of folds involving edges in G_{r-1} .

(c) $\theta : G^2 \rightarrow G_r$ is a homeomorphism.

Remark 4.8. A CT satisfies the conclusions of Theorem 2.19. This is immediate from the definitions and from Lemma 4.22.

Remark 4.9. It is an immediate consequence of (Vertices), Remark 2.8 and the definitions that a vertex whose link contains edges in more than one irreducible stratum is principal.

Remark 4.10. The tree replacement move introduced in step 2 of the proof of Theorem 2.19 shows that no taken path in zero strata is more intrinsically an edge than any other. This explains why we do not distinguish between such paths in Definition 4.4.

Remark 4.11. Exceptional paths associated to an axis $[a]_u$ have the form $E_i w^p \bar{E}_j$ where E_i and E_j are linear edges associated to $[a]_u$ such that d_i and d_j have the same sign, where $w = w_i = w_j$ and where $p \in \mathbb{Z}$.

4.2 Hard Splittings

The first item in the following lemma establishes the uniqueness of complete splittings; the second item (see also Corollary 4.13) shows that a complete splitting is a hard splitting as defined in [5].

Lemma 4.12. *Suppose that $f : G \rightarrow G$ is a CT, that σ is a circuit or path and that $\sigma = \sigma_1 \dots \sigma_m$ is a decomposition into subpaths, each of which is either a single edge in an irreducible stratum, an indivisible Nielsen path, an exceptional path or a taken connecting path in a zero stratum. Suppose also that each turn $(\bar{\sigma}_i, \sigma_{i+1})$ is legal. Then*

- (1) $\sigma = \sigma_1 \dots \sigma_m$ is the unique complete splitting of σ .
- (2) each pre-trivial subpath τ of σ is contained in a single σ_i .
- (3) a subpath of σ that has the same height as σ and is either a fixed edge or an indivisible Nielsen path equals σ_i for some i .

Proof. Let $\tilde{\sigma} = \tilde{\sigma}_1 \dots \tilde{\sigma}_m$ be a lift of σ and let $\tilde{f} : \Gamma \rightarrow \Gamma$ be a lift of $f : G \rightarrow G$. The main step in the proof is to establish the following property.

- (4) If σ_i is not a taken connecting path in a zero stratum then for each $k \geq 0$ there exist non-trivial initial and terminal subpaths $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ of $\tilde{\sigma}_i$ such that $\tilde{f}^{-k}(\tilde{f}^k(\tilde{x})) \cap \tilde{\sigma} = \{\tilde{x}\}$ for all $\tilde{x} \in \text{int}(\tilde{\alpha}_{i,k}) \cup \text{int}(\tilde{\beta}_{i,k})$.

The proof of (4) is by double induction, first on k and then on m . The $k = 0$ case is obvious so we may assume that (4) holds for any iterate less than k .

To establish the second base case, assume that $m = 1$ or equivalently that $\sigma = \sigma_i$. If σ_i is exceptional then (4) is clear (cf. Lemma 4.1.4 of [2]). If σ_i is an indivisible

Nielsen path then (4) follows from (NEG Nielsen Paths) and Lemma 2.11(2). The remaining possibility is that σ_i is an edge E in an irreducible stratum and in this case we make use of the inductive hypothesis that (4) holds in general for any iterate less than k . The first and last terms in the complete splitting of $f(E)$ are not connecting paths in zero strata. By the inductive hypothesis there exist initial and terminal subpaths $\tilde{\alpha}'$ and $\tilde{\beta}'$ of $\tilde{f}(\tilde{E})$ such that $\tilde{f}^{-(k-1)}(\tilde{f}^{k-1}(\tilde{x})) \cap \tilde{f}(\tilde{E}) = \{\tilde{x}\}$ for all $\tilde{x} \in \text{int}(\tilde{\alpha}') \cup \text{int}(\tilde{\beta}')$. Since $\tilde{f}|_{\tilde{E}}$ is an embedding, we can pull $\tilde{\alpha}'$ and $\tilde{\beta}'$ back to initial and terminal subpaths $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ of $\tilde{E}$ that satisfy (4). This completes the $m = 1$ case.

Suppose now that (4) holds for k and σ if the complete splitting of σ has fewer than $m \geq 2$ terms. As a first case suppose that σ_1 is a taken connecting path in a zero stratum H_p . By (Zero Strata), σ_2 is an edge in a EG stratum H_r with $r > p$. Define $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ using $\tilde{\sigma}_2 \dots \tilde{\sigma}_m$ in place of $\tilde{\sigma}$. Then $\tilde{f}^k(\tilde{\sigma}_1) \cap \text{int}(\tilde{f}^k(\tilde{\alpha}_{2,k})) = \emptyset$ because $\tilde{f}^k(\tilde{\sigma}_1)$ has height $< r$ and $\tilde{f}^k(\tilde{\alpha}_{2,k})$ is an embedded path whose initial direction has height r . Since $\text{int}(\tilde{f}^k(\tilde{\alpha}_{2,k}))$ separates $\tilde{f}^k(\tilde{\sigma}_1)$ from each $\tilde{f}^k(\tilde{\beta}_{i,k})$ and from $\tilde{f}^k(\tilde{\alpha}_{i,k})$ for $i > 2$, $\tilde{f}^k(\tilde{\sigma}_1)$ is disjoint from each of these sets. This proves that each $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ satisfies (4) with respect to σ .

As a second case, suppose that σ_2 is a taken connecting path in a zero stratum H_p . By (Zero Strata), σ_1 and σ_3 (if $m \geq 3$) are edges in an EG stratum H_r with $r > p$. Define $\tilde{\alpha}_{1,k}$ and $\tilde{\beta}_{1,k}$ using $\tilde{\sigma}_1$ in place of $\tilde{\sigma}$. For $i > 2$, define $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ using $\tilde{\sigma}_2 \dots \tilde{\sigma}_m$ in place of $\tilde{\sigma}$. As in the previous case, $\text{int}(\tilde{f}^k(\tilde{\beta}_{1,k})) \cap \tilde{f}^k(\tilde{\sigma}_2) = \emptyset$ and $\text{int}(\tilde{f}^k(\tilde{\alpha}_{3,k})) \cap \tilde{f}^k(\tilde{\sigma}_2) = \emptyset$. Also as in the previous case, this implies that each $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ satisfies (4) with respect to σ .

The final case is that neither σ_1 nor σ_2 is a taken connecting path in a zero stratum. Define $\tilde{\alpha}_{1,k}$ and $\tilde{\beta}_{1,k}$ using $\tilde{\sigma}_1$ in place of $\tilde{\sigma}$. For $i > 2$, define $\tilde{\alpha}_{i,k}$ and $\tilde{\beta}_{i,k}$ using $\tilde{\sigma}_2 \dots \tilde{\sigma}_m$ in place of $\tilde{\sigma}$. Since the turn $(\tilde{\sigma}_1, \tilde{\sigma}_2)$ is legal, the interiors of $\tilde{f}^k(\tilde{\alpha}_{1,k})$ and $\tilde{f}^k(\tilde{\beta}_{2,k})$ are disjoint. The proof now concludes as in the previous two cases. This completes the induction step and so the proof of (4).

If τ is a pre-trivial path then there exists $k > 0$ such that $f_{\#}^k(\tau)$ is trivial. For each $\tilde{x} \in \tilde{\tau}$ there exists $\tilde{y} \neq \tilde{x}$ in $\tilde{\tau}$ such that $\tilde{f}^k(\tilde{x}) = \tilde{f}^k(\tilde{y})$. If σ_i is not a connecting path in a zero stratum and if τ intersects $\text{int}(\sigma_i)$ then $\tau \subset \text{int}(\sigma_i)$ by (4). Since this applies to at least one of any pair of consecutive σ_i 's we have proved (2). It follows that $\sigma = \sigma_1 \cdot \dots \cdot \sigma_m$ is a splitting, and hence a complete splitting, of σ .

Suppose that $\sigma = \sigma'_1 \cdot \dots \cdot \sigma'_q$ is also a complete splitting. If σ'_i is an exceptional path or an indivisible Nielsen path then (by inspection in the NEG case and by Lemma 2.11(2) in the EG case) the interior of σ'_i is the increasing union of pre-trivial subpaths. Item (2) implies that σ'_i is contained in some σ_j . Since σ_j is not a single edge and is not contained in a zero stratum, it must be an indivisible Nielsen path or an exceptional path. By symmetry, $\sigma'_i = \sigma_j$. The terms that are taken connecting paths in zero strata are then characterized as the maximal subpaths, in the complement of the indivisible Nielsen paths and exceptional paths, that are contained in zero strata. All remaining edges are terms in the complete splitting. This proves that complete splittings are unique and so completes the proof of (1).

A fixed edge of maximal height in σ is not contained in a taken connecting path of a zero stratum, an indivisible Nielsen path or an exceptional path in σ and so must be a term in the complete splitting of σ . An indivisible Nielsen path in σ must be contained in a single σ_i by (2). If it has maximal height then, by inspection of the four possibilities for σ_i it must be all of σ_i . This proves (3). $\square$

Corollary 4.13. *Assume that $f : G \rightarrow G$ is a CT and that $\sigma = \sigma_1 \cdot \dots \cdot \sigma_s$ is the complete splitting of a path $\sigma \subset G$. If τ is an initial segment of σ with terminal endpoint in σ_j then $\tau = \sigma_1 \cdot \dots \cdot \sigma_{j-1} \cdot \mu_j$ is a splitting where μ_j is the initial segment of σ_j that is contained in τ . In particular if τ is a non-trivial Nielsen path then σ_i is a Nielsen path for all $i \leq j$ and if σ_j is not a single fixed edge then $\mu_j = \sigma_j$.*

Proof. The main statement follows immediately from Lemma 4.12(2). The statement about Nielsen paths then follows from the fact that no proper non-trivial initial segment of a non-fixed term in a complete splitting of any path is a Nielsen path. $\square$

4.3 CT versus IRT

Theorem 5.1.5 of [2] is both the definition of, and the existence theorem for, an improved relative train track map. There are eight bulleted items in the statement of the theorem, the last seven of which should be considered the definition. For notational convenience, we refer to these as (IRT-1) through (IRT-7). In this section we discuss the extent to which a CT $f : G \rightarrow G$ satisfies these seven items. By the end of this section we will have verified that CTs satisfy all of the important properties of IRTs.

(IRT-1) is that $\mathcal{F}$ is reduced, which is part of (Filtration). The following lemma states that every CT satisfies (IRT-2).

Lemma 4.14. *If $f : G \rightarrow G$ is a CT then every periodic Nielsen path has period one.*

Proof. Each periodic Nielsen path is a concatenation of periodic edges and indivisible periodic Nielsen path. The former has period one by (Periodic Edges) and the latter has period one by (Vertices) and Lemma 3.28. $\square$

The next lemma shows that a CT satisfies most of (IRT-3). The exception is that there may be some vertices v for which $f(v)$ is not fixed.

Lemma 4.15. *If $f : G \rightarrow G$ is a CT then every vertex $v \in G$ has at least two gates. If the link of v is not contained in H_r^z for some EG stratum H_r then v is principal and hence fixed.*

Proof. If (d_1, d_2) is an illegal turn then either one of the d_i 's is the terminal end of a non-fixed NEG edge or both d_1 and d_2 belong to H_r^z for some EG stratum H_r . (Vertices), (Zero Strata) and Lemma 2.10 imply that the vertex in both of these case

has two gates. At any other vertex the number of gates equals the valence. This proves the first statement of the lemma.

It follows from (Periodic Edges) and the definition of principal vertex that if v is periodic and the link of v is not contained in a single EG stratum then v is principal. If v is not periodic then its link is contained in some H_r^z by Remark 2.8 and (Zero Strata). $\square$

The difference between (IRT-4) and the conclusion of the next lemma is that a zero stratum in a IRT can be the union of contractible components.

Lemma 4.16. *Assume that $f : G \rightarrow G$ is a CT. Then G_i has a contractible component if and only if H_i is a zero stratum.*

Proof. The if direction follows from (Zero Strata). For the only if direction we assume that H_i is not a zero stratum and prove, by induction up the filtration, that every component of G_i is non-contractible. For the base case, $H_1 = G_1$ is either EG or periodic and so is connected and not contractible, by Lemma 2.10 in the former case and (Periodic Edges) in the latter. We now consider the inductive step. If some component of G_{i-1} is contractible then H_{i-1} is a zero stratum and (Zero Strata) implies that every component of G_i is non-contractible. If every component of G_{i-1} is non-contractible then (Periodic Edges) and Lemma 2.10 complete the proof. $\square$

There are two differences between (IRT-5) and (Zero Strata). The first is that an IRT can have a vertex whose link is contained in a zero stratum but a CT cannot. The second is that the restriction of an IRT to a zero stratum is always an immersion but this need not be true for a CT. We have replaced the immersion condition with Definition 4.5(2) and the assumption that every edge in a zero strata is r -taken. The primary motivation for removing the immersion condition is that it lacks robustness. For example, it need not hold for $f^2|\sigma$. Also, since the main application of relative train track maps is in analyzing the action of the induced map $f_\#$ on paths with endpoints at vertices, it makes sense to make definitions that focus on $f_\#$ and not on f .

Corollary 4.20 below implies that a CT satisfies (IRT-7). In the definition of a CT, we have replaced a list of properties satisfied by indivisible Nielsen paths corresponding to EG strata (see the statement of Corollary 4.20) with the underlying property (EG Nielsen Paths) from which these properties were derived. One advantage of this is that it is easier to deduce additional properties as in Lemma 4.25.

Lemma 4.17. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$, that ρ is an indivisible Nielsen path of height r and that ρ and H_r satisfy the conclusions of (EG Nielsen Paths). Then the illegal turn at each indivisible Nielsen path obtained by iteratively folding ρ is proper.*

Proof. Assume the notation of (EG Nielsen Paths). Since f and f_r makes exactly the same identifications on edges in H_r , each f -illegal turn in H_r is folded by f_r . Since

f_r is defined by iteratively folding ρ , say K times, the illegal turn of ρ is the only illegal turn in H_r . Let $\rho(k)$ be the indivisible Nielsen path determined by iteratively folding $\rho = \rho(0)$ k times. By hypothesis, the fold at the illegal turn of $\rho(k)$ is proper for $k < K$. It suffices to show that this holds for all k .

Let $\rho(k) = a_{1,k}b_{1,k} \dots b_{m-1,k}a_{m,k}$ be the decomposition of $\rho(k)$ into subpaths $a_{i,k} \subset H_r$ and maximal subpaths $b_{i,k} \subset G_{r-1}$. Denote the ordered sequence of the $a_{i,k}$'s by S_k . Then S_k determines the type (partial, proper, improper) of the extended fold of $\rho(k)$ and, assuming that the fold is full, also determines S_{k+1} . Since $f|_{G_r} = \theta \circ f_{r-1} \circ f_r$, $S_K = S_0$. It follows that the sequence of S_k 's is periodic with period K and hence that the fold at the illegal turn of $\rho(k)$ is proper for all k . $\square$

Lemma 4.18. *Theorem 5.15 of [4] remains true if the hypothesis that $f : G \rightarrow G$ is stable is replaced by the hypothesis that for each EG stratum H_r there is an indivisible Nielsen path ρ of height r such that ρ and H_r satisfy the conclusions of (EG Nielsen Paths).*

Proof. The proof of Theorem 5.15 has two parts. The first is a reduction to the case that the illegal turn at each indivisible Nielsen path obtained by iteratively folding ρ is full and proper. The second is the observation that in this case the proof for the special case that $f : G \rightarrow G$ is irreducible given in Lemma 3.9 of [4] applies to the general case as well. This lemma therefore follows from Lemma 4.17. $\square$

Lemma 4.19. *Proposition 5.3.1 of [2] remains true if the hypothesis $f : G \rightarrow G$ is $\mathcal{F}$ -Nielsen minimized is replaced by the hypothesis that H_r satisfies (EG Nielsen Paths).*

Proof. The proof of Proposition 5.3.1 of [2] makes use of Lemmas 5.3.6, 5.3.7, 5.3.8 and 5.3.9 of that paper. Lemma 5.3.6 states that if $f : G \rightarrow G$ is $\mathcal{F}$ -Nielsen minimized and if ρ_r crosses every edge of H_r exactly twice then H_r satisfies (EG Nielsen Paths). The remaining three lemmas use (EG Nielsen Paths) but do not refer directly to being $\mathcal{F}$ -Nielsen minimized. $\square$

The next corollary refers to *geometric strata*; complete details can be found in Definition 5.1.4 of [2].

Corollary 4.20. *Suppose that $f : G \rightarrow G$ is a relative train track map and that (EG Nielsen Paths) holds for the EG stratum H_r . Then the following properties are satisfied.*

- eg-(i)** *There is at most one indivisible Nielsen path $\rho_r \subset G_i$ that intersects H_r non-trivially. The initial edges of ρ_r and $\bar{\rho}_r$ are distinct edges in H_r .*
- eg-(ii)** *If $\rho_r \subset G_r$ is an indivisible Nielsen path that intersects H_r non-trivially and if H_r is not geometric, then there is an edge E of H_r that ρ_r crosses exactly once.*

eg-(iii) If H_r is geometric then there is an indivisible Nielsen path $\rho_r \subset G_r$ that intersects H_r non-trivially and satisfies the following properties : (i) ρ_r is a closed path with basepoint not contained in G_{r-1} ; (ii) the circuit determined by ρ_r corresponds to the unattached peripheral curve ρ^* of S ; and (iii) the surface S is connected.

In particular, H_r satisfies the EG properties of an improved relative train track.

Proof. Theorem 5.15 of [4], which applies here by Lemma 4.18, implies that there is at most one indivisible Nielsen path $\rho_r \subset G_i$ that intersects H_r non-trivially and if such a ρ_r exists then it either crosses every edge in H_r exactly twice or crosses some edge of H_r exactly once. Lemma 5.1.7 of [2] implies that if ρ_r crosses some edge of H_r exactly once then ρ_r is not a closed path. This completes the proof of eg-(i). The remaining two items follow from Proposition 5.3.1 of [2] which applies here by Lemma 4.19. $\square$

Remark 4.21. Item eg-(ii) of Corollary 4.20 and [2, Lemma 5.1.7] imply that if H_r is an EG stratum of a CT that is not geometric and if ρ is an indivisible Nielsen path of height r then ρ has distinct endpoints.

The remaining item (IRT-6) concerns NEG strata and has three parts. The first two statements of the next lemma shows that a CT satisfies the first two parts of (IRT-6). Corollary 4.24 shows that a CT satisfies the third part of (IRT-6).

Lemma 4.22. If $f : G \rightarrow G$ is a CT and H_i is NEG then H_i is a single edge E_i . If E_i is not contained in $\text{Fix}(f)$ then there is a non-trivial closed path $u_i \subset G_{i-1}$ such that $f(E_i) = E_i \cdot u_i$. Moreover u_i forms a circuit and the turn $(u_i, \bar{u}_i)$ is legal.

Proof. If H_i consists of periodic edges then the lemma follows from (Periodic Edges). Otherwise (Rotationless), (Completely Split) and (Vertices) imply that H_i is a single edge E_i and that there is a non-trivial closed path u_i such that $f(E_i) = E_i u_i$ is completely split. To prove that $f(E_i) = E_i \cdot u_i$ we must show that the first term σ_1 in the complete splitting of $E_i u_i$ is the single edge E_i . It is obviously not contained in a zero stratum and is not a Nielsen path by (NEG Nielsen Paths). It remains to show that σ_1 is not an exceptional path and for this there is no loss in assuming that E_i is linear. In the notation of (Linear Edges), $f(E_i) = E_i w_i^{d_i}$, no initial segment of which is an exceptional path by Remark 4.11. This completes the proof that $f(E_i) = E_i \cdot u_i$.

The turn $(Df^{k-1}(\bar{u}_i), Df^k(u_i))$ is the Df^{k-1} image of the legal turn $(\bar{E}_i, u_i)$ and is therefore legal for all $k \geq 1$. Since f is rotationless and since v is principal by (Vertices), $Df^k(d)$ is independent of k for all directions d based at v and all sufficiently large k . It follows that $(Df^k(\bar{u}_i), Df^k(u_i))$ is legal for all sufficiently large k and hence that $(u_i, \bar{u}_i)$ is legal. In particular, $(u_i, \bar{u}_i)$ is non-degenerate which implies that u_i forms a circuit. $\square$

Lemma 4.23. *Suppose that E_i is the unique edge of height i in a rotationless relative train track map $f : G \rightarrow G$, that $f(E_i) = E_i \cdot u_i$ for some non-trivial closed path $u_i \subset G_{i-1}$ and that every periodic Nielsen path with height less than i has period one. Suppose further that either there are no Nielsen paths of height i or E_i is a linear edge and all Nielsen paths of height i have the form $\sigma = E_i w_i^k \bar{E}_i$ where $k \neq 0$ and where w_i is root-free and $u_i = w_i^{d_i}$ for some $d_i \neq 0$. Let $h : \Gamma_{i-1} \rightarrow \Gamma_{i-1}$ be as in Notation 3.24. Then*

- $\text{Fix}(h) = \emptyset$.
- E_i is a linear edge if and only if there is a covering translation $T : \Gamma_{i-1} \rightarrow \Gamma_{i-1}$ that commutes with h and whose axis covers u_i .

Proof. Let $\tilde{f} : \Gamma \rightarrow \Gamma$ and $\tilde{E}_i$ be as in Notation 3.24. Thus $\tilde{f}(\tilde{E}_i) = \tilde{E}_i \cdot \tilde{u}_i$ where $\tilde{u}_i \subset \Gamma_{i-1}$ is a lift of u_i and h maps the initial endpoint $\tilde{x}_1$ of $\tilde{u}_i$ to the terminal endpoint $\tilde{x}_2$ of $\tilde{u}_i$. If $\tilde{v} \in \text{Fix}(h)$ and $\tilde{\gamma}$ is the path from $\tilde{x}_1$ to $\tilde{v}$ then $\tilde{E}_i \tilde{\gamma}$ is a Nielsen path for $\tilde{f}$. But then $E_i \gamma$ is a Nielsen path of height i for f that is not of the form $E_i w_i^k \bar{E}_i$. This contradiction verifies the first item.

If u_i is a Nielsen path and $T : \Gamma_{i-1} \rightarrow \Gamma_{i-1}$ is the covering translation that maps $\tilde{x}_1$ to $\tilde{x}_2$, then $Th(\tilde{x}_1) = hT(\tilde{x}_1)$ is the terminal endpoint of the lift of u_i that begins at $\tilde{x}_2$. Thus T commutes with h . For the converse suppose that h commutes with some covering translation $T : \Gamma_{i-1} \rightarrow \Gamma_{i-1}$. Corollary 3.17 implies that h is not a principal lift of $f|_{G_{i-1}}$ and hence that the endpoints of the axis of T are the only fixed points in $\partial\Gamma_{i-1}$. On the other hand, the ray $\tilde{u}_i \cdot h_{\#}(\tilde{u}_i) \cdot h_{\#}^2(\tilde{u}_i) \dots$ converges to a fixed point in $\partial\Gamma_{i-1}$. The end of this ray is therefore contained in the axis of T . It follows that u_i is a periodic Nielsen path and hence a Nielsen path and that the axis of T covers u_i . $\square$

Recall (Definition 4.1.3 of [2]) that if H_i is a NEG strata with unique edge E_i then paths of the form $E_i \gamma \bar{E}_i, E_i \gamma$ or $\gamma \bar{E}_i$ where $\gamma \subset G_{i-1}$ are called *basic paths of height i* .

Corollary 4.24. *Suppose that $f : G \rightarrow G$ is a CT and that H_i is a NEG strata with unique edge E_i . If $\sigma \subset G_i$ is a basic path of height i that does not split as a concatenation of two basic paths of height i or as a concatenation of a basic path of height i with a path contained in G_{i-1} , then either : (i) some $f_{\#}^k(\sigma)$ splits into pieces, one of which equals E_i or $\bar{E}_i$; or (ii) u_i is a Nielsen path and some $f_{\#}^k(\sigma)$ is an exceptional path of height i .*

Proof. Lemma 4.23 and Corollary 4.13 imply that $f : G \rightarrow G$ satisfies the conclusions of Proposition 5.4.3 of [2]. In conjunction with Corollary 4.20, Lemma 4.22 and Lemma 3.28 we see that $f : G \rightarrow G$ satisfies the hypotheses of Proposition 5.5.1 of [2], from which the corollary follows. $\square$

We conclude this subsection with two additional properties of CTs.

Lemma 4.25. *Suppose that $f : G \rightarrow G$ is a rotationless relative train track map, that H_r is an EG stratum satisfying (EG Nielsen Paths), and that ρ is an indivisible Nielsen path of height r . Then*

- (1) $H_r^z = H_r$.
- (2) *If $\rho = a_1 b_1 \dots b_l a_{l+1}$ is the decomposition into subpaths a_i of height r and maximal subpaths b_i of height less than r then each b_i is a Nielsen path.*
- (3) *If E is an edge of H_r then each maximal subpath of $f(E)$ in G_{r-1} is one of the b_i 's from (2). In particular $f(E)$ splits into edges in H_r and Nielsen paths in G_{r-1} .*

Proof. The maps f_r, f_{r-1} and θ induce bijections on the set of components in the filtration element of height $r - 1$. It follows that $f = \theta f_{r-1} f_r$ induces a bijection on the set of components of G_{r-1} and hence that each component of G_{r-1} is non-wandering. This proves (1).

For (2), let $(f_r)_\#(\rho) = a'_1 b'_1 \dots a'_m b'_m a'_{m+1}$ be the decomposition into subpaths a'_j of height r and maximal subpaths b'_j of height less than r . It is an immediate consequence (see the proof of Lemma 5.3.3 of [2]) of the definition of an extended fold that the set of distinct b'_j 's is contained in the set of distinct b_i 's. Now let $(\theta f_{r-1})_\#(a'_1 b'_1 \dots a'_m b'_m a'_{m+1}) = c_1 d_1 \dots c_p d_p c_{p+1}$ be the decomposition into subpaths c_k of height r and maximal subpaths d_k of height less than r . Then for each k there exists j such that $d_k = (\theta f_{r-1})_\#(b'_j)$. Combining this with the fact that $a_1 b_1 \dots b_l a_{l+1} = c_1 d_1 \dots c_p d_p c_{p+1}$, we conclude that $f_\#$ permutes the b_i 's. Since f is rotationless, each b_i is a Nielsen path.

If E is an edge of H_r then, by construction, each maximal subpath of $f_r(E)$ in G_{r-1} is a b_i . By (2), each b_i is a Nielsen path for f and hence for θf_{r-1} . This completes the proof of (3). □

Lemma 4.26. *If $f : G \rightarrow G$ is a CT and $\sigma \subset G_r$ is a path with endpoints at vertices then $f_\#^k(\sigma)$ is completely split for all sufficiently large k .*

Proof. The proof is by induction on the height of σ . The height zero case is vacuously true so suppose that σ has height $j \geq 1$ and that the lemma holds for all paths of height less than j . By Lemmas 4.6, 4.12 and the inductive hypothesis, it suffices to show that some $f^k(\sigma)$ has a splitting into subpaths that are either completely split or contained in G_{j-1} . This is immediate if H_j is a zero stratum or if H_j is a single fixed edge. If H_j is NEG then σ has a splitting into basic paths of height j and subpaths in G_{j-1} by Lemma 4.1.4 of [2]. The desired splitting of σ therefore follows from Lemma 4.24. If H_j is EG, then Lemmas 4.2.6 and 4.2.5 of [2] imply that some $f^k(\sigma)$ splits into pieces, each of which is either j -legal or a Nielsen path and Lemma 4.2.1 of [2] implies that the j -legal paths in G_j split into single edges in H_j and subpaths in G_{j-1} . □

4.4 A new move

We make use of a move that plays the same role for zero and EG strata that sliding (section 5.4 of [2]) does for NEG strata. See item (7) of Lemma 4.28 below for its main application.

Definition 4.27. Suppose that $f : G \rightarrow G$ is a rotationless relative train track map satisfying the conclusions of Theorem 2.19 with respect to the filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$, that $1 \leq j \leq N$, that every component of G_j is non-contractible and that f fixes every vertex in G_j whose link is not contained in G_j . Define a homotopy equivalence $g : G \rightarrow G$ by $g|_{G_j} = f|_{G_j}$ and $g|(G \setminus G_j) = \text{identity}$.

Define G' from G by changing the marking via g . More precisely, if X is the underlying graph of G and $\tau : R_n \rightarrow X$ is the marking that defines G , then $g\tau : R_n \rightarrow X$ is the marking that defines G' . Since G and G' have the same underlying graph, there is a natural identification of G with G' and we use this when discussing edges and strata.

Define $f' : G' \rightarrow G'$ by $f'|_{G'_j} = f|_{G_j}$ and $f'(E) = (gf)_\#(E)$ for all edges E in H_i with $i > j$.

We say that $f' : G' \rightarrow G'$ is obtained from $f : G \rightarrow G$ by *changing the marking on G_j via f* .

The following lemma is the analog of Lemma 5.4.1 of [2].

Lemma 4.28. *Suppose that $f' : G' \rightarrow G'$ is obtained from $f : G \rightarrow G$ by changing the marking on G_j via f . Then :*

- (1) $f'|_{G_j} = f|_{G_j}$.
- (2) for every path $\sigma \subset G$ with endpoints at vertices and for every $k > 0$, $g_\# f_\#^k(\sigma) = (f')_\# g_\#(\sigma)$.
- (3) $f' : G' \rightarrow G'$ is a homotopy equivalence that determines the same element of $\text{Out}(F_n)$ as $f : G \rightarrow G$.
- (4) there is a one-to-one correspondence between Nielsen paths for f and Nielsen paths for f' .
- (5) $f' : G' \rightarrow G'$ is a rotationless relative train track map satisfying the conclusions of Theorem 2.19 with respect to $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$.

Proof. Item (1) is immediate from the definitions as is the fact that f' preserves the filtration $\emptyset = G_0 \subset G_1 \subset \dots \subset G_N = G$. Also immediate are:

- (6) If $f(x) \neq f'(x)$ then $x \notin G_j$ and $f(x), f'(x) \in G_j$. In particular, $\text{Fix}(f) = \text{Fix}(f') \subset \text{Fix}(g)$, $\text{Per}(f) = \text{Per}(f') \subset \text{Per}(g)$ and Df and Df' have the same fixed and periodic directions.

- (7) Suppose that E is an edge in H_i for $i > j$ and that $f(E) = \mu_1\nu_1\mu_2\ldots\nu_{k-1}\mu_k$ where the ν_l 's are the maximal subpaths in G_j and where μ_1 and μ_k may be trivial. Then $f'(E) = \mu_1f_{\#}(\nu_1)\mu_2\ldots f_{\#}(\nu_{k-1})\mu_k$ and all the $f_{\#}(\nu_l)$'s are non-trivial. (The non-triviality follows from the fact that f fixes the endpoints of each ν_l .)

which implies

- (8) each stratum H_i has the same type (EG, NEG, zero) for f as for f' .

To verify (2), it suffices to assume that $k = 1$ and that σ is a single edge E . If $E \subset G_j$ then $g_{\#}f_{\#}(E) = f_{\#}(f_{\#}(E)) = f'_{\#}g_{\#}(E)$. If $E \subset G_i$ for $i > j$ then $g_{\#}f_{\#}(E) = f'_{\#}(E) = f'_{\#}g_{\#}(E)$. This completes the proof of (2) which implies (3).

If $\rho' \subset G'$ is a path in G with endpoints $P_1, P_2 \in \text{Fix}(f') = \text{Fix}(f)$, then there is a unique path $\rho \subset G$ with endpoints P_1 and P_2 such that $g_{\#}(\rho) = \rho'$. Condition (2) implies that ρ' is fixed by $(f')_{\#}$ if and only if ρ is fixed by $f_{\#}$. This proves (4).

To show that $f' : G' \rightarrow G'$ is a relative train track map it suffices by (1), (6) and (7) to prove that if H_i is an EG stratum with $i > j$ and if $\sigma \subset G_{i-1}$ is a connecting path for H_i then $(f')_{\#}(\sigma)$ is non-trivial. If σ is contained in a non-contractible component of G_{i-1} then its endpoints are f -fixed, and hence f' -fixed, by Remark 2.8. Non-triviality of $(f')_{\#}(\sigma)$ therefore follows from the fact that f' is a homotopy equivalence. If σ is contained in a contractible component of G_{i-1} then it is contained in a zero stratum in H_i^z so $(f')_{\#}(\sigma) = g_{\#}f_{\#}(\sigma)$. If the connecting path $f_{\#}(\sigma)$ is contained in a non-contractible component of G_{i-1} then $g_{\#}f_{\#}(\sigma)$ is non-trivial by the same argument used in the previous case. Otherwise, $g_{\#}f_{\#}(\sigma) = f_{\#}(\sigma)$ and again we are done. This completes the proof that $f' : G' \rightarrow G'$ is a relative train track map.

Item (5) follows from (4) and (6). □

4.5 Existence Theorem

Theorem 4.29. *Suppose that $\phi \in \text{Out}(F_n)$ is forward rotationless and that $\mathcal{C}$ is a nested sequence of ϕ -invariant free factor systems. Then ϕ is represented by a CT $f : G \rightarrow G$ and filtration $\mathcal{F}$ that realizes $\mathcal{C}$.*

Proof. We assume without loss $\mathcal{C}$ is maximal with respect to $\sqsubset$. Thus any filtration that realizes $\mathcal{C}$ is reduced. By Theorem 2.19 and item (3) of Lemma 2.20 we may choose a relative train track map $f : G \rightarrow G$ that represent ϕ and realizes $\mathcal{C}$ and such that each contractible component of a filtration element is a union of zero strata and the endpoints of all indivisible Nielsen paths of EG height are vertices. For the remainder of the proof all relative train track maps are assumed to satisfy these properties.

Step 1: (EG Nielsen Paths) Let $N(f)$ be the number of indivisible Nielsen paths of EG height. In the construction of an IRT in [2] it is assumed (see Definition 5.2.1

of [2]) that $N(f)$ is as small as possible. The EG properties of an IRT are then established by contradiction: the failure of these properties allows one to reduce $N(f)$ which is impossible. In order to make our constructions more algorithmic, we drop the assumption that $N(f)$ is minimal and argue inductively: the failure of (EG Nielsen Paths) allows one to reduce $N(f)$ and since $N(f)$ is finite, this process eventually terminates in an $f : G \rightarrow G$ satisfying (EG Nielsen Paths). As we are no longer assuming that $N(f)$ is minimal we cannot quote statements of results from [2] but must instead refer to their proofs.

Lemma 4.30. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ and that ρ is an indivisible Nielsen path of height r . If the fold at the illegal turn of ρ is partial then there is a relative train track map $f' : G' \rightarrow G'$ satisfying $N(f') < N(f)$.*

Proof. This follows from the proofs of Lemmas 5.2.3 and 5.2.4 of [2]. The latter constructs a topological representative of ϕ with all the desired properties and the former shows that this topological representative can be made into a relative train track map without losing any of the desired properties. $\square$

Lemma 4.31. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$, that ρ is an indivisible Nielsen path of height r and that the fold at the illegal turn of ρ is proper. Let $f' : G' \rightarrow G'$ be the relative train track map obtained from $f : G \rightarrow G$ by folding ρ . Then $N(f') = N(f)$ and there is a bijection $H_s \rightarrow H'_s$ between the EG strata of f and the EG strata of f' such that H'_s and H_s have the same number of edges for all s .*

Proof. This follows from the definition of $f' : G' \rightarrow G'$ and the proof of Lemma 5.3.3 of [2]. $\square$

Lemma 4.32. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ and that ρ is an indivisible Nielsen path of height r . If the fold at the illegal turn of ρ is improper then there is an relative train track map $f' : G' \rightarrow G'$ and a bijection $H_s \rightarrow H'_s$ between the EG strata of f and the EG strata of f' with the following properties.*

- (1) $N(f') = N(f)$.
- (2) H'_r has fewer edges than H_r .
- (3) If $s > r$ then H'_s and H_s have the same number of edges.

Proof. This follows from Definition 5.3.4 and the proof of Lemma 5.3.5 of [2]. $\square$

Lemma 4.33. *If H_r is an EG stratum of $f : G \rightarrow G$ and ρ is an indivisible Nielsen path of height r such that the illegal turn at each indivisible Nielsen path obtained by iteratively folding ρ is proper then H_r satisfies (EG Nielsen Paths).*

Proof. The conclusion of Lemma 5.3.6 of [2] is that H_r satisfies (EG Nielsen Paths). The proof of that lemma uses only standard folding arguments, the hypotheses of our lemma and uniqueness of the illegal turn of height r , which follows from Lemma 4.18. $\square$

Corollary 4.34. *Suppose that H_r is an EG stratum of a relative train track map $f : G \rightarrow G$ and that ρ is an indivisible Nielsen path of height r . Then the illegal turn at each indivisible Nielsen path obtained by iteratively folding ρ is proper if and only if H_r satisfies (EG Nielsen Paths).*

Proof. This is an immediate corollary of Lemmas 4.17 and 4.33. $\square$

Our algorithm for modifying a given $f : G \rightarrow G$ so that it satisfies (EG Nielsen Paths) is as follows. If some EG stratum does not satisfy (EG Nielsen Paths), let H_r be the highest such stratum. By Lemma 4.33, there is a (possibly empty) sequence of proper folds leading to a relative train track map and indivisible Nielsen path with either a partial fold or an improper fold. Apply Lemma 4.30 or Lemma 4.32 respectively. If the resulting relative train track map does not satisfy (EG Nielsen Paths) go back to the beginning and start again.

Remark 4.35. Iteratively folding any ρ in H_r either determines f_r as in (EG Nielsen Paths) or leads to a partial or improper fold in a predictable number of steps.

Suppose that the algorithm does not terminate. Denote the relative train track maps that are produced by $f = f_0, f_1, f_2, \dots$. Since $N(f_i)$ is non-decreasing and is strictly decreasing when a partial fold occurs, there are only finitely many such occurrences and we may assume without loss that all the folds are full. We make use of the bijection $H_s(i) \rightarrow H_s(j)$ between EG stratum for f_i and EG stratum of f_j given by Lemmas 4.31 and 4.32. Let H_r be the highest stratum for which (EG Nielsen Paths) is not satisfied by f_k for all sufficiently large k . Then the number of edges of height r is a non-increasing function of k that strictly decreases when an improper fold of height r occurs. These folds do not therefore occur for sufficiently large k . But this contradicts Lemma 4.33 and the choice of r . This proves that the algorithm terminates at a relative train track map (still called) $f : G \rightarrow G$ satisfying (EG Nielsen Paths).

In the steps that follow the number of edges in each EG strata and the number of indivisible Nielsen paths of EG height are not increased. If after some modification, (EG Nielsen Paths) fails then we can return to step 1 and start again. By the above argument this terminates after finitely many repetitions. (In fact, it is never necessary to return to step 1 but this requires an additional argument.)

Step 2: (Theorem 2.19) Apply steps (1) through (6) of the proof of Theorem 2.19 to produce a new $f : G \rightarrow G$ satisfying the conclusions of that Theorem. As noted above, we may assume by Remark 2.21 that (EG Nielsen Paths) is still satisfied.

Step 3: ((Rotationless), (Filtration) and (Zero Strata)) Items (Rotationless) and (Filtration) follow from Proposition 3.29 and Theorem 2.19(F). To achieve (Zero Strata) it suffices, by item (Z) of Theorem 2.19, to arrange that every edge in a zero stratum H_i is r -taken. Each edge E in H_i is contained in an r -taken path $\sigma \subset H_i$. If E is not r -taken, replace E by a path that has the same endpoints as σ and is marked by σ . After finitely many such tree replacements, (Zero Strata) is satisfied.

Step 4: (Periodic Edges) Suppose at first that no component C of $\text{Per}(f)$ is topologically a circle with each point in C having exactly two periodic directions. Then the endpoints of any periodic edge are principal, each periodic edge is fixed and each periodic stratum H_r has a single edge E_r . If G_{r-1} is not a core graph that contains both endpoints of E_r then one could collapse E_r without changing the free factor systems realized by the filtration elements, in violation of item (P) of Theorem 2.19). Thus (Periodic Edges) is satisfied.

For the general case, it suffices to assume that some component C of $\text{Per}(f)$ is topologically a circle with each point in C having exactly two periodic directions and modify $f : G \rightarrow G$ to reduce the number of such components.

Lemma 3.30-(1) implies that C is f -invariant and that $g = f|_C$ is orientation preserving. By (Zero Strata) and the fact that there are no periodic directions based in C and pointing out of C , every edge E_j not in C that has an endpoint in C is non-periodic, NEG and intersects C in exactly its terminal endpoint. Since all non-periodic vertices are contained in EG strata, no vertex in the complement of C maps into C . Also, C is a component of some G_l by item (NEG) of Theorem 2.19. Let E_m be the first non-periodic NEG edge E_j that has terminal endpoint in C and note that $f(E_m) = E_m C^d$. We modify $f : G \rightarrow G$ near C in two steps as follows.

In the first step we make $C \subset \text{Fix}(f)$. Extend the rotation $g^{-1} : C \rightarrow C$ to a map $h : G \rightarrow G$ that has support on a small neighborhood of C , that is homotopic to the identity and such that $h(E_j) \subset E_j \cup C$ for each non-periodic NEG edge E_j that has terminal endpoint in C . Redefine f on each edge E to be $h_{\#} f_{\#}(E)$. The filtration is unchanged. Edges in C are now fixed. If $f(E_j) = E_j u_j$ then the new u_j and the old u_j agree with the possible exception of initial and terminal segments in C . The f -image of all other edges is unchanged. In fact, $f_{\#}(\sigma)$ is unchanged for any path σ with the property that endpoints of $f(\sigma)$ are not in the support of h . It is straightforward to check that $f : G \rightarrow G$ is a relative train track map and that all of the properties that we have established to date are preserved with the possible exception of item (P) of Theorem 2.19, which fails if one or more of the E_j 's is now a fixed edge that should be collapsed. If there is no such edge then proceed to the next paragraph. If there is such an edge, collapse it as in step 3 of the proof of Theorem 2.19. That step is described very explicitly and we leave it to the reader, here and later in the proof, to check that this operation does not undo previously established properties. After finitely many such collapses, we have $C \subset \text{Fix}(f)$ and all previously established properties are preserved. If C now has outward pointing periodic directions we have finished our modifications of C . Otherwise proceed to the next paragraph.

Recall that if E_m is the first non-periodic NEG edge E_j that has terminal endpoint in C , then $f(E_m) = E_m C^d$ for some $d \in \mathbb{Z}$. In this second step we arrange that $d = 0$. Choose $h' : G \rightarrow G$ that is the identity on C , that satisfies $h'(E_j) = E_j C^{-d}$ for all E_j and that has support in a small neighborhood of C . This map is homotopic to the identity since we can simply unwind the twisting on C . Redefine f on each edge E to be $h_{\#} f_{\#}(E)$ and note that $C \cup E_m \subset \text{Fix}(f)$ so the component of $\text{Fix}(f)$ containing C is no longer a topological circle. The filtration is unchanged. If necessary, collapse fixed edges with an endpoint in C and repeat this second step.

Step 5: (Induction: the NEG case) It remains to establish (Completely Split) and the items related to non-fixed NEG edges. We do this by induction up the filtration making use of sliding and the new move described in Section 4.4.

Let NI be the number of irreducible strata in the filtration and for each $0 \leq m \leq NI$, let $G_{i(m)}$ be the smallest filtration element containing the first m irreducible strata. We will prove by induction on m that for all $0 \leq m \leq NI$, one can modify f to arrange that $f|_{G_{i(m)}}$ is a CT. The $m = 0$ case is vacuously true so we assume that $f|_{G_r}$ is a CT for $r = i(m)$ and make modifications to arrange that $f|_{G_s}$ is a CT for $s = i(m + 1)$. In this step we assume that H_s is NEG and is hence a single edge E_s satisfying $f(E_s) = E_s u_s$ for some path $u_s \subset G_{s-1}$.

By (Zero Strata), $r = s - 1$. The sliding operation described in section 2.7 (complete details in section 5.4 of [2]) allows us to modify E_s and u_s by choosing a path $\tau \subset G_{s-1}$ with initial endpoint equal to the terminal vertex of E_s and ‘sliding’ the terminal end of E_s to the terminal vertex of τ . As noted in step 2, we may assume that sliding preserves (EG Nielsen Paths).

As a first case suppose that after sliding along τ we have $E_s \subset \text{Fix}(f)$. For future reference note that by Lemma 2.17 this is equivalent to $[\bar{\tau} u_s f_{\#}(\tau)]$ being trivial and hence equivalent to $f_{\#}(E_s \tau) = E_s [u_s f_{\#}(\tau)] = E_s \tau$; i.e., to $E_s \tau$ being a Nielsen path.

If both endpoints of E_s are contained in G_{s-1} then (Periodic Edges) is satisfied as are all of the conclusions of Theorem 2.19 and the three properties established in step 3. The remaining properties of a CT follow from the inductive hypothesis.

If either endpoint of E_s is not contained in G_{s-1} then collapse E_s to a point as in step 4. None of the previously achieved properties are lost and the remaining properties of a CT follow from the inductive hypothesis. This completes the inductive step in the case that $E_s \subset \text{Fix}(f)$ is trivial after sliding.

We assume now that there is no choice of τ such that $E_s \tau$ is a Nielsen path. The following proposition is a combination of Propositions 5.4.3 and 5.5.1 of [2].

Proposition 4.36. *Suppose that*

- (i) $f : G \rightarrow G$ is a relative train track map that satisfies (EG Nielsen Paths)
- (ii) $f|_{G_{s-1}}$ satisfies (Completely Split)
- (iii) H_s is an NEG stratum with single edge E_s for which there does not exist a path $\mu \subset G_{s-1}$ such that $E_s \mu$ is a Nielsen path.

Then there exists a path $\tau \subset G_{s-1}$ with initial endpoint equal to the terminal endpoint of E_s such after performing the slide associated to τ the following conditions are satisfied.

- (1) $f(E_s) = E_s \cdot u_s$ is a non-trivial splitting.
- (2) If σ is a circuit or path with endpoints at vertices and if σ has height s then there exists $k \geq 0$ such that $f_{\#}^k(\sigma)$ splits into subpaths of the following type.
 - (a) E_s or $\bar{E}_s$
 - (b) an exceptional path of height s
 - (c) a subpath of G_{s-1}
- (3) u_s is completely split and its initial vertex is principal.
- (4) $f|G_s$ satisfies (Linear Edges).

Proof. The construction of a path τ along which to slide is carried out in the proof of Proposition 5.4.3 of [2]. We assume that τ has been chosen to satisfy the conclusions of that proposition. In particular, $f(E_s) = E_s \cdot u_s$ is a splitting that is non-trivial by (iii). Thus (1) is satisfied. (The statement of Proposition 5.4.3 of [2] allows the possibility that G is subdivided at a periodic point and that the terminal endpoint of E_s is one of the new periodic vertices. By the end of the construction, we will have shown that the terminal endpoint of E_s is principal and hence fixed. At that point we can undo the subdivision.)

For (3) we must make use of facts that are explicitly stated and used in the proof of Proposition 5.4.3 of [2] but are not contained in its statement. The first is that by a further slide one can replace u_s with $f_{\#}^k(u_s)$ for any $k \geq 1$. Since $f|G_{s-1}$ satisfies (Completely Split) we may assume by Lemma 4.26 that u_s is completely split. The second is that if $u_s = \alpha \cdot \beta$ is a coarsening of the complete splitting of u_s , then by a further slide we may assume that the terminal endpoint of (the new) E_s is the terminal endpoint of α . Thus to complete the proof of (3) we need only show that the endpoint of some term in the complete splitting of u_s is principal. The only way that this could fail would be if u_s has height r' where $H_{r'}$ is EG and if each height r' term in the complete splitting of u_s is a single edge. After replacing u_s with a sufficiently high iterate, we may assume that u_s has such a long r' -legal segment that every edge in $H_{r'}$ occurs as a term in the complete splitting of u_s . Lemma 3.19 then completes the proof of (3).

If E_s is a linear edge, choose a root-free Nielsen path w_s and $d_s \neq 0$ so that $u_s = w_s^{d_s}$. If $E_t \subset G_{s-1}$ is a linear edge with the same axis as E_s then after reversing the orientation on w_s we may assume that w_t and w_s agree as oriented loops. After a further slide as in the proof of (3) we may assume that $w_s = w_t$. Item (iii) implies that $E_s \bar{E}_t$ is not a Nielsen path and hence that $d_s \neq d_t$. This completes the proof of item (4).

Lemma 4.1.4 of [2] states that if σ is a height s circuit or path with endpoints at vertices then σ splits into subpaths that are either contained in G_{s-1} or are basic paths of height s meaning that they, or their inverse has the form $E_s\gamma$ or $E_s\gamma\bar{E}_s$ for some $\gamma \subset G_{s-1}$. By Corollary 4.20, $f : G \rightarrow G$ satisfies the hypotheses, and hence the conclusions, of Proposition 5.5.1 of [2]. These conclusions address both types of basic paths of height s and verify (2). $\square$

We assume now that we have performed the slide move of Proposition 4.36. Since u_s is non-trivial, $f|_{G_s}$ satisfies (Periodic Edges) and all of the properties achieved in the first four steps of our construction. Items (Completely Split), (Vertices), (NEG Strata), (Linear Edges) and (Nielsen Paths) for $f|_{G_s}$ follows from Proposition 4.36 and these properties for $f|_{G_{s-1}}$. This completes the proof of the inductive step in the case that H_s is NEG.

Step 6: (Induction: the EG case) Suppose now that H_s is EG. items (Vertices), (NEG Strata), (Linear Edges) and (Nielsen Paths) for $f|_{G_s}$ follows from these properties for $f|_{G_{s-1}}$.

For each edge $E \subset H_s$, there is a decomposition $f(E) = \mu_1 \cdot \nu_1 \cdot \mu_2 \cdot \dots \cdot \nu_{m-1} \cdot \mu_m$ where the ν_l 's are the maximal subpaths in G_r . Let $\{\nu_l\}$ be the collection of all such paths that occur as E varies over the edges of H_s . By (RTT-ii), $f_{\#}^k(\nu_l)$ is non-trivial for each k and l . By Lemma 4.26 we may choose k so large that each $f_{\#}^k(\nu_l)$ is completely split. We may also assume that the endpoints of $f_{\#}^k(\nu_l)$ are periodic and hence principal. There are finitely many connecting paths σ contained in the strata (if any) between G_r and H_s . Each $f(\sigma)$ is either a connecting path or a non-trivial path in G_r with fixed endpoints. We may therefore assume that $f_{\#}^k(\sigma)$ is completely split for each such σ . After k applications of Lemma 4.28 with $j = r$ (see in particular item (7) of that lemma) we have that $f|_{G_s}$ is completely split. This completes the induction step and so also the proof of the theorem. $\square$

4.6 Further properties of a CT

The following lemma is an extension of Lemma 3.26.

Lemma 4.37. *Assume that $f : G \rightarrow G$ is a CT. The following properties hold for every principal lift $\tilde{f} : \Gamma \rightarrow \Gamma$.*

- (1) *If $\tilde{v} \in \text{Fix}(\tilde{f})$ and a non-fixed edge $\tilde{E}$ determines a fixed direction at $\tilde{v}$, then $\tilde{E} \subset \tilde{f}_{\#}(\tilde{E}) \subset \tilde{f}_{\#}^2(\tilde{E}) \subset \dots$ is an increasing sequence of paths whose union is a ray $\tilde{R}$ that converges to some $P \in \text{Fix}_N(\hat{f})$ and whose interior is fixed point free.*
- (2) *For every isolated $P \in \text{Fix}_N(\hat{f})$ there exists $\tilde{E}$ and $\tilde{R}$ as in (1) that converges to P . The edge E is non-linear.*

Proof. For $\tilde{E}$ as in (1) and for each $m > 0$, Lemma 4.6 implies that $\tilde{E} \subset \tilde{f}_\#(\tilde{E}) \subset \tilde{f}_\#^2(\tilde{E}) \subset \dots$ is a nested sequence of completely split paths. This increasing sequence of paths defines a ray $\tilde{R}'$ that converges to some non-repelling fixed point $P \in \text{Fix}_N(\hat{f})$ and that, by Corollary 4.13, intersects $\text{Fix}(\hat{f})$ only in its initial endpoint. This completes the proof of (1).

If $P \in \text{Fix}_N(\hat{f})$ is isolated then $\tilde{f}$ moves points that are sufficiently close to P toward P by Lemma 2.3. We may therefore choose a ray $\tilde{R}$ that converges to P and that intersects $\text{Fix}(\hat{f})$ only in its initial endpoint. Moreover, the initial edge $\tilde{E}$ of $\tilde{R}$ determines a fixed direction by Lemma 3.16 and so extends to a fixed point free ray $\tilde{R}$ converging to some $Q \in \text{Fix}_N(\hat{f})$ by (1). Lemma 3.16 implies that $P = Q$. Since P is isolated it is not an endpoint of the axis of a covering translation and E is not a linear edge. $\square$

Notation 4.38. If $\tilde{E}$ and P are as in Lemma 4.37(1) then we say that $\tilde{E}$ *iterates to* P and that P is *associated to* $\tilde{E}$.

The following lemma is used in section 5 and also in [10].

Lemma 4.39. *Suppose that $\psi \in \text{Out}(F_n)$ is forward rotationless and that $P \in \text{Fix}_N(\hat{\Psi})$ for some $\Psi \in P(\psi)$. Suppose further that Λ is an attracting lamination for some element of $\text{Out}(F_n)$, that Λ is ψ -invariant and that Λ is contained in the accumulation set of P . Then $\text{PF}_\Lambda(\psi) \geq 0$ and $\text{PF}_\Lambda(\psi) > 0$ if and only if P is isolated in $\text{Fix}_N(\hat{\Psi})$.*

Proof. Let $g : G \rightarrow G$ be a CT representing ψ , let $\tilde{g} : \Gamma \rightarrow \Gamma$ be the lift corresponding to Ψ and let $\tilde{R} \subset \Gamma$ be a ray converging to P . Choose a generic leaf $\gamma \subset G$ of the realization of Λ in G . Then every finite subpath of γ lifts to a subpath of every subray of $\tilde{R}$. If P is not isolated in $\text{Fix}_N(\hat{\Psi})$ then Lemma 2.3 implies that every finite subpath of γ extends to a Nielsen path for g . By (EG Strata) and (Nielsen Paths), this extension can be done with a uniformly bounded number of edges. It is an immediate consequence of the definition of the expansion factor (Definition 3.3.2 of [2]) that $\text{PF}_\Lambda(\psi) = 0$.

Assume now that P is isolated in $\text{Fix}_N(\hat{\Psi})$ and let Σ be the set of finite paths $\sigma \subset G$ with endpoints at vertices and with the property that every finite subpath of γ is contained in $g_\#^m(\sigma)$ for some $m > 0$. Lemma 4.37 and the assumption that Λ is contained in the accumulation set of P imply that Σ contains a path that is a single edge and in particular is non-empty.

Let $\sigma \in \Sigma$ be an element of minimal height, say k . Then σ decomposes as a concatenation of edges $\mu_i \subset H_k$ and subpaths $\nu_i \subset G_{k-1}$ and we let K be the number of elements in this decomposition. Choose a nested sequence of subpaths γ_j of γ whose union equals γ . Since the ν_i 's are not in Σ , there exists $J > 0$ so that γ_j is not contained in any $g_\#^m(\nu_i)$ for $j > J$. Since γ is generic, it is birecurrent. Choose $j_0 > J$ and $j > j_0$ so that γ_j contains at least $3K$ disjoint copies of γ_{j_0} . There exists $m > 0$

such that $\gamma_j \subset g_{\#}^m(\sigma)$. It follows that $\gamma_{j_0} \subset g_{\#}^m(\mu_i)$ for some i . There is a choice of i that works for all choices of j_0 and this proves that $\mu_i \in \Sigma$.

Let E be the single edge in μ_i . We assume that E is NEG and argue to a contradiction. There is a path $u \subset G_{k-1}$ such that $g_{\#}^m(E) = E \cdot u \cdot \dots \cdot g_{\#}^{m-1}(u)$ for all $m > 0$. Lemma 3.1.16 of [2] states that γ is not a circuit. It follows that u is not a Nielsen path and hence that the length of $g_{\#}^l(u)$ goes to infinity with l . The birecurrence of γ and the fact that $E \in \Sigma$ imply that for every γ_j there exists $p > 0$ such that $\gamma_j \subset g_{\#}^{p-1}(u)g_{\#}^p(u) = g_{\#}^{p-1}(ug_{\#}(u))$ in contradiction to the assumption that no element of Σ has height less than k .

We now know that H_k is EG. Since $E \in \Sigma$, γ is a leaf in the attracting lamination Λ_k associated to H_k . There is a splitting of $g(E)$ into subpaths in H_k and subpaths in G_{k-1} . If γ were contained in G_{k-1} then one of the subpaths in G_{k-1} would be contained in Σ in contradiction to our choice of k . Thus γ is not entirely contained in G_{k-1} and Lemma 3.1.15 of [2] implies that γ is a generic leaf of Λ_k . In other words, $\Lambda = \Lambda_k$. It follows that $\text{PF}_{\Lambda}(\psi) > 0$ which completes the proof. $\square$

Assume that ϕ is forward rotationless and that $f : G \rightarrow G$ is a CT representing ϕ . Following the notation of [3] we say that an unoriented conjugacy class μ of a root-free element of F_n is an *axis for ϕ* if for some (and hence any) representative $c \in F_n$ there exist distinct $\Phi_1, \Phi_2 \in \text{P}(\phi)$ that fix c . Equivalently $\text{Fix}_N(\hat{\Phi}_1) \cap \text{Fix}_N(\hat{\Phi}_2)$ is the endpoint set of the axis A_c for T_c . It is a consequence of Lemma 4.41 below that an unoriented conjugacy class μ is an axis for ϕ if and only if it is an axis for a linear edge in some (every) CT representing ϕ .

Remark 4.40. In the context of the mapping class group, a conjugacy class is an axis if and only if it is represented by a reducing curve in the minimal Thurston normal form.

Lemma 3.30 implies that the oriented conjugacy class of c is ϕ -invariant. By Lemmas 4.1.4 and 4.2.6 of [2], the circuit γ representing c splits into a concatenation of subpaths α_i , each of which is either a fixed edge or an indivisible Nielsen path. (NEG Nielsen Paths) and Corollary 4.20 imply that each turn $(\bar{\alpha}_i, \alpha_{i+1})$ is legal. Item 1 of Lemma 4.12 therefore implies that this splitting is the complete splitting of γ .

There is an induced complete splitting of A_c into subpaths $\tilde{\alpha}_i$ that project to either fixed edges or indivisible Nielsen paths. The lift $\tilde{f}_0 : \Gamma \rightarrow \Gamma$ that fixes the endpoints of each $\tilde{\alpha}_i$ is a principal lift by Corollary 3.27 and commutes with T_c . We say that $\tilde{f}_0$ and the corresponding $\Phi_0 \in \text{P}(\phi)$ are the *base lift* and *base principal automorphism* associated to μ and the choices of T_c and $f : G \rightarrow G$. (If μ is not represented by a basis element then Φ_0 is independent of the choice of $f : G \rightarrow G$ but otherwise it is not. Indeed, if μ is not represented by a basis element, then the smallest free factor F that carries μ has rank greater than one. Since μ is not an axis for $\phi|F$ there is a unique principal automorphism $\Phi|F$ that fixes c and Φ must be the extension of $\Phi|F$.) Item 2 of Lemma 4.12 implies that for each $\tilde{\alpha}_i$ and for each $\tilde{x} \in \tilde{\alpha}_i$, the nearest

point to $\tilde{f}_0(\tilde{x})$ in A_c is contained in $\tilde{\alpha}_i$. It follows that $\text{Fix}(T_c^j \tilde{f}_0) = \emptyset$ for all $j \neq 0$ and hence that $\tilde{f}_0$ is the only lift that commutes with T_c and has fixed points in A_c .

Lemma 4.41. *Suppose that ϕ is forward rotationless and that the unoriented conjugacy class μ is an axis for ϕ . Assume notation as above. There is a bijection between the set of principal lifts [principal automorphisms] $\tilde{f}_j \neq \tilde{f}_0$ [respectively $\Phi_j \neq \Phi_0 \in P(\phi)$] that commute with T_c [fix c] and the set of linear edges $\{E_j\}$ with axis equal to μ . Moreover, if $f(E_j) = E_j w_j^{d_j}$ then $\tilde{f}_j = T_c^{d_j} \tilde{f}_0$ [$\Phi_j = i_c^{d_j} \Phi_0$].*

Proof. The w_j 's in question are equal by (Linear Edges) and we label this path w . There is a lift $\tilde{w}$ that is a fundamental domain of A_c and that is a Nielsen path for $\tilde{f}_0$. Let $\tilde{E}_j$ be the lift of E_j that terminates at the initial endpoint $\tilde{v}$ of $\tilde{w}$ and let $\tilde{f}_j$ be the lift that fixes the initial endpoint of $\tilde{E}_j$. Then $\tilde{f}_j$ is a principal lift that commutes with T_c and satisfies $\tilde{f}_j(\tilde{v}) = T_c^{d_j}(\tilde{v}) = T_c^{d_j}(\tilde{f}_0 \tilde{v})$ which implies that $\tilde{f}_j = T_c^{d_j} \tilde{f}_0$.

Conversely, if $\tilde{f} \neq \tilde{f}_0$ is a principal lift that commutes with T_c then $\tilde{f} = T_c^d \tilde{f}_0$ for some $d \neq 0$. In particular, A_c is disjoint from $\text{Fix}(\tilde{f})$ and there is a ray $\tilde{R}_1$ that intersects $\text{Fix}(\tilde{f})$ in exactly its initial endpoint and that terminates at the endpoint P of A_c that is the limit of the forward $\tilde{f}$ orbit of $\tilde{v}$. Let $\tilde{E}$ be the initial edge of $\tilde{R}_1$. Lemma 3.16 implies that $\tilde{E}$ determines a fixed direction and also that $\tilde{R}_1$ must be the ray constructed from the initial edge $\tilde{E}$ of $\tilde{R}_1$ by Lemma 4.37. If $\tilde{f}(\tilde{E}) = \tilde{E} \cdot \tilde{u}$ then $\tilde{R}_1 = \tilde{E} \cdot \tilde{u} \cdot \tilde{f}_\#(\tilde{u}) \cdot \tilde{f}_\#^2(\tilde{u}) \cdot \dots$. Since $\tilde{R}_1$ has a common infinite end with A_c , it follows that $\tilde{f}_\#^k(u)$ is a periodic, hence fixed, Nielsen path for sufficiently large k and for u equal to the projected image of $\tilde{u}$. In particular, u and $\tilde{f}_\#(u)$ have the same $\tilde{f}_\#^k$ -image, and since they have the same endpoints, they must be equal. In other words, u is a Nielsen path. This proves that $\tilde{E}$ is the lift of a linear edge E whose associated axis is μ . By (Linear Edges), $E = E_j$ and $d = d_j$ for some j and, after translating $\tilde{E}$ by some iterate of T_c if necessary, that $\tilde{v}$ is the terminal endpoint of $\tilde{E}$. $\square$

Remark 4.42. Suppose that $f : G \rightarrow G$ is a CT, that C is a component of some filtration element G_s , that C has no valence one vertices and that $\phi|_{\mathcal{F}(C)}$ is the trivial outer automorphism. Then $f|_C$ is the identity. To see this, let H_r be the first non-fixed stratum in C . It can not be EG because the identity element has no attracting laminations. If it were NEG it would have to be linear because $f|_{G_{r-1}}$ is the identity and it cannot be linear because the identity element has no axes.

We conclude this section by showing that every element of $\text{Out}(F_n)$ has a uniformly bounded iterate that is forward rotationless.

Lemma 4.43. *For all $n \geq 1$ there exists $K_n > 1$ so that ϕ^{K_n} is forward rotationless for all $\phi \in \text{Out}(F_n)$.*

Proof. Given $\phi \in \text{Out}(F_n)$, let $f : G \rightarrow G$ be a CT representing some forward rotationless iterate $\psi = \phi^N$ of ϕ . By Corollary 3.17 and Lemma 3.8, the number of isogredience classes of principal lifts of ψ is the same as the number of Nielsen classes

for $f : G \rightarrow G$. If x is a principal vertex that has valence two and that is isolated in $\text{Fix}(f)$ then x is either the initial endpoint of a non-fixed NEG edge or an endpoint of an indivisible Nielsen path of EG height. By (Vertices) and Corollary 4.20, there is a uniform (i.e. depending only on n) upper bound to the number of isolated fixed principal vertices. By (Periodic Edges) there is also a uniform upper bound to the number of components of $\text{Fix}(f)$ that contain at least one edge. It follows that there is a uniform upper bound to the number of Nielsen classes for $f : G \rightarrow G$ and to the number of edges based at principal vertices. From the former we conclude that the number of isogredience classes of principal lifts $\tilde{f} : \Gamma \rightarrow \Gamma$ of f is uniformly bounded.

Since ϕ commutes with ψ , it acts on the set of isogredience classes of principal automorphisms representing ψ . After replacing ϕ with a uniformly bounded iterate, we may assume that ϕ fixes each isogredience class. Thus, if Ψ is a principal automorphism representing ψ then there exists an automorphism Φ representing ϕ such that Φ commutes with Ψ . In particular, $\mathbb{F} := \text{Fix}(\Psi)$ is Φ -invariant. By construction, the outer automorphism determined by $\Phi|_{\mathbb{F}}$ has finite order and so is represented by a homeomorphism of a graph with no valence one or valence two vertices. Since the rank of $\mathbb{F}$ is uniformly bounded, the period of the outer automorphism determined by $\Phi|_{\mathbb{F}}$ is uniformly bounded. After replacing ϕ with a further uniformly bounded iterate, we may assume that $\mathbb{F} \subset \text{Fix}(\Phi)$. Thus $\text{Fix}(\hat{\Phi})$ contains each non-isolated point of $\text{Fix}(\hat{\Psi})$ by Lemma 2.3.

By Lemma 4.37(2), the number of isolated points in $\text{Fix}(\hat{\Psi})$, up to the action of $\mathbb{F}$, is bounded above by the number of edges based at principal fixed points for f and so is uniformly bounded. We may therefore assume that if P is an isolated point in $\text{Fix}(\hat{\Psi})$ then $\hat{\Phi}(P) = \hat{T}_a(P)$ for some $a = a_P \in \mathbb{F}$, from which it follows that $\hat{\Phi}^N(P) = \hat{T}_a^N(P)$.

The proof now divides into cases. If $\mathbb{F}$ is trivial then $\text{Fix}(\hat{\Psi}) \subset \text{Fix}(\hat{\Phi})$. If $\mathbb{F}$ has rank at least two then $\Phi^N = \Psi$. It follows that T_a is trivial and again $\text{Fix}(\hat{\Psi}) \subset \text{Fix}(\hat{\Phi})$. The final case is that $\mathbb{F}$ has rank one. After replacing Φ with $T_a^{-1}\Phi$ we may assume that $P \in \text{Fix}(\hat{\Phi})$. Since $\text{Fix}(\hat{\Phi})$ and $\text{Fix}(\hat{\Psi})$ have at least three points in common, $\Phi^N = \Psi$. As in the higher rank case, it follows that $\text{Fix}(\hat{\Psi}) \subset \text{Fix}(\hat{\Phi})$ in this case as well. As this holds for each principal automorphism representing ψ , ϕ is forward rotationless. $\square$

5 Recognition Theorem

In this section we specify invariants that uniquely determine a forward rotationless ϕ . As a warm-up to the general theorem, we consider the special case, essentially proved in [1], that ϕ is irreducible, meaning that there are no non-trivial proper ϕ -invariant free factor systems. It follows that a CT $f : G \rightarrow G$ representing ϕ has a single stratum and that the stratum is EG. In particular, ϕ has infinite order and $\mathcal{L}(\phi)$ has exactly one element. Lemma 3.30(3) implies that all iterates of ϕ are irreducible.

Lemma 5.1. *If $\phi \in \text{Out}(F_n)$ is irreducible and forward rotationless, then ϕ has infinite order and is determined by its unique attracting lamination Λ and the expansion factor $\text{PF}_\Lambda(\phi)$. More precisely, if ϕ and ψ are forward rotationless and irreducible and if they have the same unique attracting lamination and the same expansion factor then $\phi = \psi$.*

Proof. As noted above, ϕ and ψ have infinite order and all iterates of ϕ and ψ are irreducible. Theorem 2.14 of [1] implies that $\psi^{-1}\phi$ has finite order and that $\psi^k = \phi^k$ for some $k \geq 1$. By Lemma 3.19 and Lemma 2.13 there exists $\Phi \in P(\phi)$ such that $\text{Fix}_N(\Phi)$ contains at least three points P_1, P_2 and P_3 , each of whose accumulation set equals Λ . The Fix_N -preserving bijections between $P(\phi)$ and $P(\phi^k)$ and between $P(\psi)$ and $P(\psi^k)$ induce a Fix_N -preserving bijection between $P(\phi)$ and $P(\psi)$. Thus there exists $\Psi \in P(\psi)$ such that $\text{Fix}_N(\hat{\Psi}) = \text{Fix}_N(\hat{\Phi})$.

Choose a finite order homeomorphism $f : G \rightarrow G$ of a marked graph G representing $\psi^{-1}\phi$, let $\tilde{f} : \Gamma \rightarrow \Gamma$ be the lift corresponding to $\Psi^{-1}\Phi$ and note that $P_1, P_2, P_3 \in \text{Fix}(\tilde{f})$. The line with endpoints P_1 and P_2 and the line with endpoints P_1 and P_3 are $\tilde{f}_\#$ -invariant and since $\tilde{f}$ is a homeomorphism they are $\tilde{f}$ -invariant. The intersection of these lines is an $\tilde{f}$ -invariant, and hence $\tilde{f}$ -fixed, ray $\tilde{R}$ that terminates at P_1 . The lamination Λ is carried by the subgraph $G_0 \subset \text{Fix}(f)$ that is the image of $\tilde{R}$. Example 2.5(1) of [1] implies that $G_0 = G$; thus f is the identity and $\psi = \phi$. $\square$

If $\Phi_1, \Phi_2 \in P(\phi)$ and if there exists a non-trivial indivisible $a \in \text{Fix}(\Phi_1) \cap \text{Fix}(\Phi_2)$, then $\Phi_2 \Phi_1^{-1} = i_a^d$ for some $d \neq 0$. We think of d as a *twist coefficient* for the ordered pair (Φ_1, Φ_2) relative to a . In our next example we show that an elementary linear outer automorphism is determined by the fixed subgroups of its principal automorphisms and by a twist coefficient.

Example 5.2. Let $x_1, x_2, \dots, x_n$ be a basis for F_n and let $F_{n-1} = \langle x_1, \dots, x_{n-1} \rangle$. Define Φ_1 by $\Phi_1|_{F_{n-1}} = \text{identity}$ and $\Phi_1(x_n) = x_n a^d$ for some non-trivial root-free $a \in F_{n-1}$ and some $d > 0$. Define

$$\Phi_2 = i_{x_n}^{-1} \Phi_1 i_{x_n} = i_{\bar{x}_n \Phi_1(x_n)} \Phi_1 = i_a^d \Phi_1.$$

Then $\text{Fix}(\Phi_2) = i_{\bar{x}_n} \text{Fix}(\Phi_1)$ and $\text{Fix}(\Phi_1) \cap \text{Fix}(\Phi_2) = \langle a \rangle$. Since $\text{Fix}(\Phi_1)$ and $\text{Fix}(\Phi_2)$ have rank greater than one, $\Phi_1, \Phi_2 \in P(\phi)$.

We claim that if there exist $\Psi_1, \Psi_2 \in P(\psi)$ such that $\Psi_2 = i_a^d \Psi_1$ and such that $\text{Fix}(\Psi_i) = \text{Fix}(\Phi_i)$ for $i = 1, 2$, then $\psi = \phi$. It is obvious that $\Psi_1|_{F_{n-1}} = \text{identity}$. Moreover,

$$\begin{aligned} \text{Fix}(i_a^d \Psi_1) &= \text{Fix}(i_a^d \Phi_1) = \text{Fix}(i_{x_n}^{-1} \Phi_1 i_{x_n}) = i_{\bar{x}_n} \text{Fix}(\Phi_1) \\ &= i_{\bar{x}_n} \text{Fix}(\Psi_1) = \text{Fix}(i_{x_n}^{-1} \Psi_1 i_{x_n}) = \text{Fix}(i_{\bar{x}_n \Psi_1(x_n)} \Psi_1). \end{aligned}$$

Since $i_a^d \Psi_1$ and $i_{\bar{x}_n \Psi_1(x_n)} \Psi_1$ represent the same outer automorphism and have a common fixed subgroup of rank greater than one, they are equal. Thus $a^d = \bar{x}_n \Psi_1(x_n)$ or equivalently $\Psi_1(x_n) = x_n a^d$. This proves that $\Psi_1 = \Phi_1$ and $\phi = \psi$.

We now turn to the general case.

Theorem 5.3. (Recognition Theorem) *Suppose that $\phi, \psi \in \text{Out}(F_n)$ are forward rotationless and that*

(1) $\text{PF}_\Lambda(\phi) = \text{PF}_\Lambda(\psi)$, for all $\Lambda \in \mathcal{L}(\phi) = \mathcal{L}(\psi)$.

(2) *there is bijection $B : P(\phi) \rightarrow P(\psi)$ such that:*

(i) **(fixed sets preserved)** $\text{Fix}_N(\hat{\Phi}) = \text{Fix}_N(\widehat{B(\Phi)})$

(ii) **(twist coordinates preserved)** *If $w \in \text{Fix}(\Phi)$ and $\Phi, i_w \Phi \in P(\phi)$, then $B(i_w \Phi) = i_w B(\Phi)$.*

Then $\phi = \psi$.

Remark 5.4. The bijection B is necessarily equivariant in the sense that $B(i_c \Phi i_c^{-1}) = i_c B(\Phi) i_c^{-1}$ for all $c \in F_n$. This follows from the fact that $\text{Fix}(i_c \Phi i_c^{-1}) = i_c(\text{Fix}(\Phi))$ and from Remark 3.4. Thus B is determined by its value on one representative from each of the finitely many equivalence classes in $P(\phi)$ and 2(i) can be verified by checking finitely many cases. Similarly, 2(ii) can be verified by checking finitely many cases. The w 's to which 2(ii) apply have the form $w = a^d$ where a represents a common axis of ϕ and ψ . The values of d can be read off from relative train track maps as in Lemma 4.41.

Remark 5.5. The assumption in (1) that $\mathcal{L}(\phi) = \mathcal{L}(\psi)$ is redundant. It follows from Lemma 3.26 and 2(i). We include it in the statement of the theorem for clarity.

Proof. The proof is by induction on n . By convention, all forward rotationless outer automorphisms are the identity when $n = 1$ so we may assume that the theorem holds for all ranks less than n and prove it for n .

The case that both ϕ and ψ are irreducible is proved in Lemma 5.1 so we may assume that at least one of these, say ϕ , is reducible and so admits a proper non-trivial invariant free factor system. Since this free factor system is realized by a filtration element in a relative train track map representing ϕ , some proper free factor carries either an attracting lamination Λ for ϕ or a ϕ -periodic conjugacy class $[c]$. Lemma 3.30 implies that the elements of $\mathcal{L}(\phi) = \mathcal{L}(\psi)$ are invariant by both ϕ and ψ , that ϕ and ψ have the same periodic conjugacy classes and that all these conjugacy classes are fixed. The smallest free factor that carries Λ or $[c]$ is both ϕ -invariant and ψ -invariant by Corollary 2.5. This proves the existence of non-trivial proper free factors that are that both ϕ -invariant and ψ -invariant.

Among all proper free factor systems, each of whose elements is both ϕ -invariant and ψ -invariant, choose one $\mathcal{F} = \{[F^1], \dots, [F^k]\}$ that is maximal with respect to inclusion. We claim that $\phi|F^i = \psi|F^i$ for each i . If F^i has rank one then this follows from Lemma 3.30. If F^i has rank at least two then principal automorphisms representing $\phi|F^i$ and $\psi|F^i$ extend uniquely to principal automorphisms representing

ϕ and ψ . Thus $\phi|F^i$ and $\psi|F^i$, which are forward rotationless by Corollary 3.31, satisfy the hypothesis of Theorem 5.3 and the inductive hypothesis implies that $\phi|F^i = \psi|F^i$. This verifies the claim. Let $f : G \rightarrow G$ be a CT representing ϕ with $[\pi_1(G_r)] = \mathcal{F}$ for some filtration element G_r , which we may assume without loss has no valence one vertices. Then $f|G_r$ represents both $\phi|\mathcal{F}$ and $\psi|\mathcal{F}$. The proof now divides into two cases. The arguments are sufficiently elaborate that we treat the cases in separate subsections.

5.1 The NEG Case

In this subsection we complete the proof of Theorem 5.3 in the case that there exists $s > r$ such that G_s is not homotopy equivalent to G_r and such that H_i is NEG for all $r < i \leq s$. After reordering the H_i 's if necessary, we may assume that G_s is obtained from G_r as a topological space by either adding a disjoint circle or by attaching an arc E with both endpoints in G_r . In the former case, $[\pi_1(G_s)]$ is both ϕ -invariant and ψ -invariant in contradiction to the assumption that $\mathcal{F}$ is maximal and the fact that G_s is disconnected. Thus $G_s = G_r \cup E$ where $f(E) = \bar{u}_1 E u_2$ for some closed paths $u_1, u_2 \subset G_r$. If E is a single edge of G then $s = r + 1$ and at least one of u_1 or u_2 is trivial. Otherwise E is made up of two edges and $s = r + 2$.

Choose lifts $\tilde{E}, \tilde{u}_1, \tilde{u}_2 \subset \Gamma$ and $\tilde{f} : \Gamma \rightarrow \Gamma$ such that $\tilde{f}(\tilde{E}) = \tilde{u}_1^{-1} \tilde{E} \tilde{u}_2$. Denote the component of G_r that contains u_i by C^i and the copy of the universal cover of C^i that contains $\tilde{u}_i$ by Γ_r^i . If u_1 or u_2 is trivial then (Remark 4.9) at least one of the endpoints of $\tilde{E}$ is a principal vertex that is fixed by $\tilde{f}$. Otherwise $\tilde{E}$ subdivides into two NEG edges whose common initial vertex is principal and is fixed by $\tilde{f}$. Corollary 3.27 therefore implies that $\tilde{f}$ is a principal lift. Lemma 3.26 implies that there is a line $\tilde{\gamma} \subset \Gamma$ that crosses $\tilde{E}$ and has endpoints in $\text{Fix}_N(\tilde{f})$. The smallest free factor system that carries $[\pi_1(G_r)]$ and γ is both ϕ -invariant and ψ -invariant. Since $\mathcal{F}$ is maximal, $G = G_s$.

Corollary 3.2.2 of [2] implies that ψ is represented by $g : G \rightarrow G$ such that $g|G_r = f|G_r$ and such that $g(E) = \bar{w}_1 E w_2$ for some closed paths $w_1, w_2 \subset G_r$. It suffices to prove that $u_i = w_i$. The cases are symmetric so we show that $u_1 = w_1$.

Suppose at first that C^1 has rank one and hence is a topological circle that is contained in $\text{Fix}(f)$. By (Periodic Edges), the vertices in C^1 are principal. Thus at least one of E or $\bar{E}$ determines a direction based in C^1 that is fixed by Df . If u_1 is non-trivial then it must be that $\bar{E}$ determines a fixed direction based in C^1 . In this case $C^1 \cup E$ is a component of G and hence equal to G . We conclude that $n = 2$ and that there is a basis $\{x_1, x_2\}$ for F_2 and $d \neq 0$ such that $x_1 \mapsto x_1$ and $x_2 \mapsto x_2 x_2^d$ defines an automorphism representing ϕ . This is a special case of Example 5.2 and so $u_1 = w_1$ in this case. We may therefore assume that u_1 is trivial. The symmetric argument with g replacing f reduces us to the case that u_1 and w_1 are both trivial and so equal. We may now assume that C^1 has rank at least two.

The principal lift $\tilde{g} : \Gamma \rightarrow \Gamma$ that corresponds to $\tilde{f}$ under the bijection B satisfies

$\text{Fix}_N(\hat{g}) = \text{Fix}_N(\hat{f})$. Since $\hat{g}$ fixes the endpoints of $\tilde{\gamma}$ and $\tilde{E}$ is the only edge in $\tilde{\gamma}$ that does not project into G_r , it follows that $\tilde{g}(\tilde{E}) = \tilde{w}_1 \tilde{E} \tilde{w}_2$. Let $\tilde{v} \in \Gamma_r^1$ be the initial endpoint of $\tilde{E}$. Then $\tilde{u}_1$ and $\tilde{w}_1$ are the paths in Γ_r^1 connecting $\tilde{v}$ to $\tilde{f}(\tilde{v})$ and $\tilde{v}$ to $\tilde{g}(\tilde{v})$ respectively. It therefore suffices to show that $\tilde{f}|_{\Gamma_r^1} = \tilde{g}|_{\Gamma_r^1}$.

We know that $\tilde{f}|_{\Gamma_r^1}$ and $\tilde{g}|_{\Gamma_r^1}$ are both lifts of $f|_{C^1}$ and that $\hat{f}|_{\partial\Gamma_r^1}$ and $\hat{g}|_{\partial\Gamma_r^1}$ have a common fixed point P . If P is not an endpoint of the axis of some covering translation T_c of Γ_r^1 , then there is at most one lift of $f|_{C^1}$ that fixes P and we are done. Suppose then that $P \in \{T_c^\pm\}$. By Remark 3.2, there exists a principal lift of $f|_{C^1}$ that commutes with T_c . This lift extends over Γ to principal lifts $\tilde{f}'$ and $\tilde{g}'$ of f and g respectively. Since $\text{Fix}_N(\hat{f}|_{\partial\Gamma_{r-1}}) \subset \text{Fix}_N(\tilde{f}') \cap \text{Fix}_N(\tilde{g}')$ contains at least three points, $\tilde{g}' = B(\tilde{f}')$. Condition 2(ii) therefore implies that $\tilde{g} = T_c^d \tilde{g}'$ and $\tilde{f} = T_c^d \tilde{f}'$ for some $d \in \mathbb{Z}$. We conclude that $\tilde{f}|_{\Gamma_r^1} = \tilde{g}|_{\Gamma_r^1}$ as desired.

5.2 The EG Case

In this subsection we prove Theorem 5.3 assuming that there exists $s > r$ where H_s is exponentially growing and where the union of the non-contractible components of G_{s-1} is homotopy equivalent to G_r . In light of subsection 5.1, this completes the proof of Theorem 5.3. Since G_{s-1} and G_r carry the same elements of $\mathcal{L}(\phi)$, all irreducible strata between G_r and G_s are NEG. Our choice of $\mathcal{F}$ guarantees that $G_s = G$.

Denote $\psi^{-1}\phi$ by θ . We must show that θ is trivial. By construction, $\theta[[F^l]]$ is trivial for each $[[F^l]] \in \mathcal{F}$ and the attracting lamination Λ associated to H_s is θ -invariant with expansion factor one. Moreover, for any principal lift Φ of ϕ there is a unique lift Θ of θ such that $\text{Fix}(\hat{\Theta}) \supset \text{Fix}_N(\hat{\Phi})$.

Each F^l corresponds to a non-contractible component D_l of G_{s-1} . Let $\tilde{D}_l$ be the component of the full pre-image of D_l whose accumulation set in ∂F_n is ∂F^l . Suppose that $v \in H_s \cap D_l$ and that $\tilde{v} \in \tilde{D}_l$ is a lift of v . Then $\tilde{v}$ is principal by Remark 4.9 and the lift $\tilde{f}_{\tilde{v}}$ of f that fixes $\tilde{v}$ is principal by Lemma 3.27. The link of $\tilde{v}$ contains edges that project to H_s and determine fixed directions for $\tilde{f}_{\tilde{v}}$. Lemma 3.26 implies that any such edge extends to a fixed point free ray that terminates at a point $P \in \text{Fix}_N(\hat{f}_{\tilde{v}})$ whose accumulation set is Λ . Let $\mathcal{P}_l$ be the union of such P for all $v \in H_s \cap D_l$ and all lifts $\tilde{v} \in \tilde{D}_l$.

Lemma 5.6. *There is a lift Θ of θ such that $\partial F^l \cup \mathcal{P}_l \subset \text{Fix}(\hat{\Theta})$.*

Proof. Assume at first that F^l has rank at least two. Let Θ be the unique lift of θ such that $\partial F^l \subset \text{Fix}(\hat{\Theta})$. If $P \in \mathcal{P}_l$ corresponds to $\tilde{v}$ as above and if $\Phi_1 \in \mathcal{P}(\phi)$ corresponds to $\tilde{f}_{\tilde{v}}$ then $\text{Fix}_N(\hat{\Phi}_1)$ contains P and intersects ∂F^l non-trivially. There exists Θ_1 such that $\text{Fix}(\hat{\Theta}_1) \supset \text{Fix}_N(\hat{\Phi}_1)$. If $\text{Fix}(\hat{\Phi}_1) \cap \partial F^l \neq \{T_c^\pm\}$ for some covering translation T_c , then $\Theta_1 = \Theta$ and we are done. Suppose then that $\text{Fix}(\hat{\Phi}_1) \cap \partial F^l = T_c^\pm$. By Remark 3.2 there is a principal lift Φ_2 such that $\text{Fix}(\Phi_2)$ contains $T_c^\pm$ and such that $\Phi_2|_{F_l}$ is a principal lift of $\phi|_{F_l}$. In particular, $\Phi_2 = i_c^d \Phi_1$ for some $d \neq 0$. By hypothesis, there are principal lifts Ψ_1 and Ψ_2 such that $\text{Fix}_N(\hat{\Psi}_i) = \text{Fix}_N(\hat{\Phi}_i)$ and

such that $\Psi_2 = i_c^d \Psi_1$. Thus $\Theta = \Psi_2^{-1} \Phi_2 = \Psi_1^{-1} \Phi_1 = \Theta_1$ where the first equality comes from the fact that $\text{Fix}(\hat{\Psi}_2^{-1} \hat{\Phi}_2)$ contains at least three points in ∂F^l .

It remains to consider the case that F^l has rank one. For each $P \in \mathcal{P}_l$, there exist Φ_P and Ψ_P such that $\text{Fix}_N(\hat{\Phi}_P) = \text{Fix}_N(\hat{\Psi}_P)$ contains $P \cup \partial F^l$. Define $\Theta_P = \Psi_P^{-1} \Phi_P$. For any $Q \in \mathcal{P}_l$ there exists w such that $\Phi_P = i_w \Phi_Q$ and $\Psi_P = i_w \Psi_Q$. It follows that $\Theta_P = \Psi_P^{-1} \Phi_P = \Psi_Q^{-1} \Phi_Q = \Theta_Q$ as desired. $\square$

Corollary 5.7. *If θ has finite order then θ is trivial.*

Proof. If θ has finite order then [8] there is a marked graph X , a subgraph X_0 such that $\mathcal{F}(X_0) = \mathcal{F}$ and a homeomorphism $h : X \rightarrow X$ that represents θ and is the identity on X_0 . By Lemma 5.6 there is an $h_\#$ -invariant R whose initial endpoint is in X_0 and whose accumulation set contains Λ . No proper free factor system carries $\mathcal{F}$ and Λ , so R crosses every edge in $X \setminus X_0$. Since h is a homeomorphism $R \subset \text{Fix}(h)$ and we conclude that h is the identity. $\square$

We now assume that θ has infinite order and argue to a contradiction. There is no loss in replacing θ by an iterate, so we may assume that both θ and θ^{-1} are forward rotationless. There is a CT $h : G' \rightarrow G'$ representing θ and there exists $r' < s'$ such that $G'_{s'} = G'$, such that $\mathcal{F}(G'_{r'}) = \mathcal{F}$ and such that $h|_{G'_{r'}} = \text{identity}$ (see Remark 4.42).

Lemma 5.8. *Suppose that $\mathcal{P}_l$ and Θ are as in Lemma 5.6 and that $P \in \mathcal{P}_l$. Then P is not isolated in $\text{Fix}(\hat{\Theta})$.*

Proof. Suppose at first that P is an attractor for $\hat{\Theta}$. Let $\tilde{h}$ be the lift of h corresponding to Θ . By Lemma 4.37 there is an edge $\tilde{E}$ that iterates to P ; let $\tilde{R}$ be the ray connecting $\tilde{E}$ to P . If E belongs to an EG stratum, then Λ , which is the accumulation set of P , is an attracting lamination for θ by Lemma 4.39. This contradicts the fact that θ acts on Λ with expansion factor one. If E is NEG, then Λ is carried by $G' \setminus E$ in contradiction to the fact that no proper free factor can carry $\mathcal{F}$ and Λ . This proves that P is not an attractor for $\hat{\Theta}$.

The symmetric argument using a relative train track map for θ^{-1} proves that P is not a repeller so Lemma 2.3 completes the proof. $\square$

Corollary 5.9. *If $\gamma' \subset G'$ is a finite subpath of either:*

- (1) *a leaf of the realization of Λ in G' or*
- (2) *the projection of the line in Γ' connecting a pair of points $P_1, P_2 \in \mathcal{P}_l$*

then γ' extends to a Nielsen path for h .

Proof. Let Θ be as in Lemma 5.6 and let $\tilde{h} : \Gamma' \rightarrow \Gamma'$ be the lift corresponding to Θ . For case (1), let $\tilde{R}' \subset \Gamma'$ be a ray converging to $P \in \mathcal{P}_l$. There are lifts $\tilde{\gamma}' \subset \tilde{R}'$ of γ' that are arbitrarily close to P . Lemma 5.8 and Lemma 2.3 therefore imply that $\tilde{\gamma}'$ extends to a Nielsen path for $\tilde{h}$. In case (2), $P_1, P_2 \in \text{Fix}(\hat{h})$. Lemma 5.8 and Lemma 2.3 imply that any finite subpath of the line connecting P_1 to P_2 extends to a Nielsen path for $\tilde{h}$. $\square$

It is well known that if ϕ acts trivially on conjugacy classes in F_n then ϕ is the trivial element. This can be proved by induction up the strata of $f : G \rightarrow G$ representing ϕ or directly as in Lemma 3.3 of [9]. The following lemma therefore completes the proof of Theorem 5.3.

Lemma 5.10. *θ fixes each conjugacy class $[c]$ in F_n .*

Proof. If v is a vertex in G whose link $lk(v)$ is contained in H_s , then the local stable Whitehead graph SW_v is defined to be the graph with one vertex for each oriented edge based at v whose initial direction is fixed by Df and an edge connecting the vertices corresponding to E_1 and E_2 if there is an edge E of H_s and $k \geq 1$ so that the path $f_{\#}^k(E)$ contains $\bar{E}_1 E_2$ or $\bar{E}_2 E_1$ as a subpath. By Lemma 2.13 this is equivalent to $\bar{E}_1 E_2$ or $\bar{E}_2 E_1$ being a subpath of a generic leaf of Λ . If SW_v is not connected then then one can blow up v to an edge E as in Proposition 5.4 of [4] to obtain a proper free factor that carries $\mathcal{F}$ and Λ . Since this is impossible, SW_v is connected.

Choose a positive integer M such that Df^M maps every direction in H_s to a fixed direction in H_s . At one point in the proof we need a way to choose partial edges and for this we subdivide the edges of H_s at the full f^M -pre-image of the set of vertices. Edges in this subdivision will be called *edgelets*. Thus an edgelet maps by f^M to an edge.

Let $g : G \rightarrow G'$ be a homotopy equivalence that respects the marking and that satisfies $g(G_r) = G'_{r'}$. We show below that there is a positive integer N so that for all circuits $\sigma \subset G$ the conjugacy class in F_n determined by $g_{\#} f_{\#}^{MN}(\sigma) \subset G'$ is fixed by θ . Since every conjugacy class in F_n is realized in this manner by some σ , this completes the proof of the lemma.

Since θ acts by the identity on $\mathcal{F}(G'_{r'})$ we may assume without loss that σ crosses at least one edge in H_s . The proof involves choosing a closed curve that is homotopic to $f_{\#}^{MN}(\sigma)$ and a covering of that curve by subpaths with large overlap.

To begin, choose a cyclic ordering of the m edges of H_s in σ . Define σ_1 to be first edge of H_s in σ , σ_3 to be second edge of H_s in σ and σ_2 to be the subpath of σ that begins with the last edgelet in σ_1 and ends with the first edgelet in σ_3 . Define σ_5 to be third edge of H_s in σ and σ_4 to be the subpath of σ that begins with the last edgelet in σ_3 and ends with the first edgelet in the σ_5 . Continue in this manner stopping with σ_{2m} that begins with the last edgelet in σ_{2m-1} and ends with the first edgelet in σ_1 .

Let $\rho = f_{\#}^M(\sigma)$ and let $\rho_i = f_{\#}^M(\sigma_i)$. Then each ρ_i is an edge path in G whose initial and terminal edges are in H_s and whose initial and terminal directions are

fixed by Df . Suppose that $\rho_{2j} = \bar{E}E'$ where the link of the common initial endpoint v of E and E' is entirely contained in H_s . Since SW_v is connected, there are edges $E = E_1, E_2, \dots, E_l = E'$ in $lk(v)$ with initial directions fixed by Df such that each $\bar{E}_p E_{p+1}$ is a subpath of a generic leaf of Λ . Replace ρ_{2j} by the concatenation $(\bar{E}_1 E_2) \cdot (\bar{E}_2 E_3) \cdot \dots \cdot (\bar{E}_{l-1} E_l)$.

After adjusting the indices, we have produced paths $\rho_1, \dots, \rho_t$ with the following properties:

- (a) The initial edge E_i^1 of ρ_i and the terminal edge E_i^2 of ρ_i are contained in H_s and E_i^2 equals E_{i+1}^1 up to a possible change of orientation.
- (b) For all $k \geq 0$, $f_{\#}^k(\rho_i)$ is a finite subpath of either:
 - (1) a generic leaf of the realization of Λ in G or
 - (2) the projection of the line in Γ connecting a pair of points $P_1, P_2 \in \mathcal{P}_l$.

Suppose that each E_i^1 as been decomposed into proper subpaths $E_i^1 = a_i b_i$. The equality $E_i^2 = E_{i+1}^1$ or $\bar{E}_{i+1}^1$ determines a corresponding decomposition of E_i^2 . Define τ_i from ρ_i by deleting the initial a_i segment of E_i^1 and by deleting the terminal segment of E_i^2 determined from (a) as follows. If $E_i^2 = E_{i+1}^1$ then remove the terminal b_{i+1} ; if $E_i^2 = \bar{E}_{i+1}^1$ then remove the terminal a_{i+1} .

- (c) For any $\{a_i\}$ as above, ρ is homotopic to the loop determined by the concatenation of the τ_i 's.

Choose K greater than the number of edges with height s' in any indivisible Nielsen path for h . By [7] there is a positive constant C so that if $\beta_1 \subset \beta_2$ are finite subpaths in G then $g_{\#}(\beta_2) \subset G'$ contains the subpath of $g_{\#}(\beta_1)$ obtained by removing the initial and terminal segments of edge length C . Since generic leaves of Λ are birecurrent and since the realization of Λ in G' can not be contained in $G'_{r'}$, there is a subpath γ' of a generic leaf of the realization of Λ in G' that contains $2K + 3C$ edges of $H'_{s'}$. Choose a subpath γ of a generic leaf of the realization of Λ in G whose $g_{\#}$ image contains γ' . There exists $N > 0$ so that $f_{\#}^N(E)$ contains γ as a subpath for each edge E of H_s . It follows that the path $g_{\#}(f_{\#}^N(E))$ contains at least $2K + C$ edges of $H'_{s'}$ for each edge E of H_s .

The subpath ν'_i of $g_{\#}(f_{\#}^N(\rho_i))$ defined by removing initial and terminal segments with exactly C edges of $H'_{s'}$ is contained in either the realization of a leaf of Λ in G' or the projection of a line in Γ' connecting a pair of points $P_1, P_2 \in \mathcal{P}_l$. Corollary 5.9 implies that ν'_i extends to a Nielsen path μ'_i for h . Let $\mu'_i = \mu'_{i,1} \cdot \mu'_{i,2} \cdot \dots \cdot \mu'_{i,m_i}$ be the complete splitting of μ'_i . There is no loss in assuming that $\mu'_{i,2} \subset \nu'_i$. There are at most $K + C$ edges of $H'_{s'}$ in $g_{\#}(f_{\#}^N(\rho_i))$ that precede $\mu'_{i,2}$. Without changing this estimate we may assume that $\mu'_{i,2}$ has height s' . Note that $\mu'_{i,2} \subset g_{\#}(f_{\#}^N(E_i^1))$ and hence that $\mu'_{i+1,2} \subset g_{\#}(f_{\#}^N(E_i^2))$. Let a_i be an initial segment of E_i^1 such that

$g_{\#}f_{\#}^N(a_i)$ is the initial segment of $g_{\#}(f_{\#}^N(\rho_i))$ that precedes $\mu'_{i,2}$. Define the τ_i 's as in (c).

Lemma 4.12(3) implies that there exists $l \leq m_i$ such that $\mu'_{i,l} = \mu'_{i+1,2}$ up to a change of orientation. Thus $g_{\#}f_{\#}^N(\tau_i) = \mu'_{i,2} \cdots \mu'_{i,l}$ is a Nielsen path for h . Property (c) implies that the conjugacy class determined by $g_{\#}f_{\#}^N(\rho) = g_{\#}f_{\#}^{MN}(\sigma)$ is θ -invariant as desired. $\square$

Corollary 5.11. *If ϕ and ψ are forward rotationless and if $\phi^m = \psi^m$ for some $m > 0$ then $\phi = \psi$.*

Proof. Since ϕ and ψ are forward rotationless there are Fix_N -preserving bijections between $P(\phi^m)$ and $P(\phi)$ and between $P(\psi^m)$ and $P(\psi)$. By assumption, $P(\phi^m) = P(\psi^m)$ so there is a Fix_N -preserving bijection between $P(\phi)$ and $P(\psi)$. The lemma now follows from the Recognition theorem and the fact that expansion factors and twist coefficients for ϕ^m are m times those of ϕ and similarly for ψ . $\square$

References

- [1] Mladen Bestvina, Mark Feighn, and Michael Handel. Laminations, trees, and irreducible automorphisms of free groups. *Geom. Funct. Anal.*, 7(2):215–244, 1997.
- [2] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for $\text{Out}(F_n)$. I. Dynamics of exponentially-growing automorphisms. *Ann. of Math. (2)*, 151(2):517–623, 2000.
- [3] Mladen Bestvina, Mark Feighn, and Michael Handel. Solvable subgroups of $\text{Out}(F_n)$ are virtually Abelian. *Geom. Dedicata*, 104:71–96, 2004.
- [4] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [5] Martin Bridson and Daniel Groves. Free-group automorphisms, train tracks and the beaded decomposition. preprint, 2005.
- [6] Marshall M. Cohen and Martin Lustig. The conjugacy problem for Dehn twist automorphisms of free groups. *Comment. Math. Helv.*, 74(2):179–200, 1999.
- [7] Daryl Cooper. Automorphisms of free groups have finitely generated fixed point sets. *J. Algebra*, 111(2):453–456, 1987.
- [8] Marc Culler. Finite groups of outer automorphisms of a free group. In *Contributions to group theory*, volume 33 of *Contemp. Math.*, pages 197–207. Amer. Math. Soc., Providence, RI, 1984.

- [9] Benson Farb and Michael Handel. Commensurations of $\text{Out}(F_n)$. *Publ. Math. Inst. Hautes Études Sci.*, (105):1–48, 2007.
- [10] Mark Feighn and Michael Handel. Abelian subgroups of $\text{Out}(F_n)$. to appear in *Geometry and Topology*.
- [11] William J. Floyd. Group completions and limit sets of Kleinian groups. *Invent. Math.*, 57(3):205–218, 1980.
- [12] Damien Gaboriau, Andre Jaeger, Gilbert Levitt, and Martin Lustig. An index for counting fixed points of automorphisms of free groups. *Duke Math. J.*, 93(3):425–452, 1998.
- [13] Michael Handel and William P. Thurston. New proofs of some results of Nielsen. *Adv. in Math.*, 56(2):173–191, 1985.
- [14] Gilbert Levitt and Martin Lustig. Periodic ends, growth rates, Hölder dynamics for automorphisms of free groups. *Comment. Math. Helv.*, 75(3):415–429, 2000.
- [15] Gilbert Levitt and Martin Lustig. Automorphisms of free groups have asymptotically periodic dynamics. *J. Reine Angew. Math.*, 619:1–36, 2008.
- [16] Richard T. Miller. Geodesic laminations from Nielsen’s viewpoint. *Adv. in Math.*, 45(2):189–212, 1982.